WORLD INTELLECTUAL PROPERTY ORGANIZA International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:

C12N 15/86, 15/35, 5/10, A61K 48/00

(11) International Publication Number:

WO 00/28061

(43) International Publication Date:

18 May 2000 (18.05.00)

(21) International Application Number:

PCT/US99/25694

A3

(22) International Filing Date:

2 November 1999 (02.11.99)

(30) Priority Data:

60/107,114

5 November 1998 (05.11.98)

US

(71) Applicant (for all designated States except US): TRUSTEES OF THE UNIVERSITY OF PENNSYLVA-NIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104-3147 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): WILSON, James, M. [US/US]; 1350 N. Avignon Drive, Gladwyne, PA 19035 (US). XIAO, Weidong [CN/US]; Apartment P4, 155 Washington Lane, Jenkintown, PA 19046 (US).
- (74) Agents: KODROFF, Cathy, A. et al.; Howson & Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(88) Date of publication of the international search report: 3 August 2000 (03.08.00)

(54) Title: ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAIN-**ING SAME**

(57) Abstract

The nucleic acid sequences of adeno-associated virus (AAV) serotype 1 are provided, as are vectors and host cells containing these sequences and functional fragments thereof. Also provided are methods of delivering genes via AAV-1 derived vectors.

Express Mail No. ET033648243US

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	AL	Albania	ES	C:-		_		
	AM.			Spain	LS	Lesotho	SI	Slovenia
		Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
	AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
i	AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ĺ	AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
ļ	BA	Bosnia and Herzegovina	GE	Georgia .	MD	Republic of Moldova	TG	Togo
	BB	Barbados	GH	Ghaпа	MG	Madagascar	TJ	Tajikistan
ı	BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
	BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
į	BC	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
į	BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
	BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
	BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
ĺ	CA	Canada	IT	Italy	MX	Mexico	UZ 、	Uzbekistan
	CF	Central African Republic	JP	Јарап	NE	Niger	VN	Viet Nam
	CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
	СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
	CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	211	Zimoabwe
1	CM	Cameroon		Republic of Korea	PL	Poland		
	CN	China	KR	Republic of Korea	PT	Portugal		•
	CU	Cuba	KZ	Kazakstan	RO	Romania		
ı	CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
	DE	Germany	LI	Liechtenstein	SD	Sudan		
ı	DK	Denmark	LK	Sri Lanka	SE			
	; EE	Estonia	LR	Liberia	SG	Sweden		
ı					30	Singapore	•	
1								

ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

This work was supported by the National Institutes of Health, grant no. P30 DK47757-06 and PO1 HD32649-04. The US government may have certain rights in this invention.

Field of the Invention

5

This invention relates generally to viral vector, and more particularly, to recombinant viral vectors useful for gene delivery.

Background of the Invention

10 Adeno-associated viruses are small, single-stranded DNA viruses which require helper virus to facilitate efficient replication [K.I. Berns, Parvoviridae: the viruses and their replication, p. 1007-1041, in F.N. Fields et al., Fundamental virology, 3rd ed., vol. 2, (Lippencott-Raven Publishers, Philadelphia, PA) (1995)]. The 4.7 kb genome of AAV is characterized by two inverted terminal repeats (ITR) and two open reading frames which encode the Rep proteins and Cap proteins, 15 respectively. The Rep reading frame encodes four proteins of molecular weight 78 kD, 68 kD, 52 kD and 40 kD. These proteins function mainly in regulating AAV replication and integration of the AAV into a host cell's chromosomes. The Cap reading frame encodes three structural proteins in molecular weight 85 kD (VP 1), 72 kD (VP2) and 61 kD (VP3) [Berns, cited above]. More than 80% of total proteins in 20 AAV virion comprise VP3. The two ITRs are the only cis elements essential for AAV replication, packaging and integration. There are two conformations of AAV ITRs called "flip" and "flop". These differences in conformation originated from the replication model of adeno-associated virus which use the ITR to initiate and reinitiate the replication [R.O. Snyder et al., J. Virol., 67:6096-6104 (1993); K.I. Berns, 25 Microbiological Reviews, 54:316-329 (1990)].

AAVs have been found in many animal species, including primates, canine, fowl and human [F.A. Murphy et al., "The Classification and Nomenclature of Viruses: Sixth Report of the International Committee on Taxonomy of Viruses",

2

Archives of Virology, (Springer-Verlag, Vienna) (1995)]. In addition to five known primate AAVs (AAV-1 to AAV-5), AAV-6, another serotype closely related to AAV-2 and AAV-1 has also been isolated [E. A. Rutledge et al., J. Virol., 72:309-319 (1998)]. Among all known AAV serotypes, AAV-2 is perhaps the most well-characterized serotype, because its infectious clone was the first made [R.J. Samulski et al., Proc. Natl. Acad. Sci. USA, 79:2077-2081 (1982)]. Subsequently, the full sequences for AAV-3A, AAV-3B, AAV-4 and AAV-6 have also been determined [Rutledge, cited above; J.A.Chiorini et al., J. Virol., 71:6823-6833 (1997), S. Muramatsu et al., Virol., 221:208-217 (1996)]. Generally, all AAVs share more than 80% homology in nucleotide sequence.

5

10

25

A number of unique properties make AAV a promising vector for human gene therapy [Muzyczka, Current Topics in Microbiology and Immunology, 158:97-129 (1992)]. Unlike other viral vectors, AAVs have not been shown to be associated with any known human disease and are generally not considered pathogenic. Wild type AAV is capable of integrating into host chromosomes in a site specific manner [R. M. 15 Kotin et al., Proc. Natl. Acad. Sci. USA, 87:2211-2215 (1990)- R.J. Samulski, EMBO J., 10(12):3941-3950 (1991)]. Recombinant AAV vectors can integrate into tissue cultured cells in chromosome 19 if the rep proteins are supplied in trans [C. Balague et al., J. Virol., 71:3299-3306 (1997); R. T. Surosky et al., J. Virol., 71:7951-7959 (1997)]. The integrated genomes of AAV have been shown to allow 20 long term gene expression in a number of tissues, including, muscle, liver, and brain [K. J. Fisher, Nature Med., 3(3):306-312 (1997); R. O. Snyder et al., Nature Genetics, 16:270-276 (1997); X. Xiao et al., Experimental Neurology, 144:113-124 (1997); Xiao, J. Virol., 70(11):8098-8108 (1996)].

AAV-2 has been shown to be present in about 80-90% of the human population. Earlier studies showed that neutralizing antibodies for AAV-2 are prevalent [W. P. Parks et al., <u>J. Virol.</u>, <u>2</u>:716-722 (1970)]. The presence of such antibodies may significantly decrease the usefulness of AAV vectors based on AAV-2 despite its other merits. What are needed in the art are vectors characterized by the

3

advantages of AAV-2, including those described above, without the disadvantages, including the presence of neutralizing antibodies.

Summary of the Invention

5

10

15

20

25

In one aspect, the invention provides an isolated AAV-1 nucleic acid molecule which is selected from among SEQ ID NO: 1, the strand complementary to SEQ ID NO: 1, and cDNA and RNA sequences complementary to SEQ ID NO: 1 and its complementary strand.

In another aspect, the present invention provides AAV ITR sequences, which include the 5' ITR sequences, nt 1 to 143 of SEQ ID NO: 1; the 3' ITR sequences, nt 4576 to 4718 of SEQ ID NO: 1, and fragments thereof.

In yet another aspect, the present invention provides a recombinant vector comprising an AAV-1 ITR and a selected transgene. Preferably, the vector comprises both the 5' and 3' AAV-1 ITRs between which the selected transgene is located.

In still another aspect, the invention provides a recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.

In a further aspect, the present invention provides a nucleic acid molecule encoding an AAV-1 rep coding region and an AAV-1 cap coding region.

In still another aspect, the present invention provides a host cell transduced with a recombinant viral vector of the invention. The invention further provides a host cell stably transduced with an AAV-1 P5 promoter of the invention.

In still a further aspect, the present invention provides a pharmaceutical composition comprising a carrier and a vector of the invention.

In yet another aspect, the present invention provides a method for AAV-mediated delivery of a transgene to a host involving the step of delivering to a selected
host a recombinant viral vector comprising a selected transgene under the control of
sequences which direct expression thereof and an adeno-associated virus 1 (AAV-1)
virion.

15

20

25

In another aspect, the invention provides a method for in vitro production of a selected gene product using a vector of the invention.

Other aspects and advantages of the invention will be readily apparent to one of skill in the art from the detailed description of the invention.

5 Brief Description of the Drawings

Figs. 1A-1C illustrate the alignment of nucleotides of AAV-1 [SEQ ID NO: 1], AAV-2 [SEQ ID NO: 18] and AAV-6 [SEQ ID NO: 19]. The alignment was done with MacVector 6.0. The full sequences of AAV-1 are shown in the top line. Nucleotides in AAV-2 and AAV-6 identical to AAV-1 are symbolized by "." and gaps by "-". Some of the conserved features among AAVs are marked in this figure. Note the 3' ITRs of AAV-1 and AAV-6 are shown in different orientations.

Fig. 2 illustrates the predicted secondary structure of AAV-1 ITR. The nucleotides in AAV-2 and AAV-6 are shown in italic and bold respectively.

Fig. 3A illustrates a hypothesis of how AAV-6 arose from the homologous recombination between AAV-1 and AAV-2. The major elements of AAV-1 are indicated in the graph. A region that is shared between AAV-1, AAV-2 and AAV-6 is shown in box with waved lines.

Fig. 3B is a detailed illustration of a 71 bp homologous region among AAV-1, AAV-2 and AAV-6. Nucleotides that differ among these serotypes are indicated by arrows.

Fig. 4A is a bar chart illustrating expression levels of human alpha 1 antitrypsin (α1AT) in serum following delivery of hAAT via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 4B is a bar chart illustrating expression levels of erythropoietin (epo) in serum following delivery of the epo gene via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 5A is a bar chart illustrating expression levels of $\alpha 1$ AT in liver following delivery of $\alpha 1$ AT as described in Example 7.

10

20

25

Fig. 5B is a bar chart demonstrating expression levels of epo in liver following delivery of epo as described in Example 7.

Fig. 5C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to liver as described in Example 7.

Fig. 5D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of alAT or epo to liver as described in Example 7.

Fig. 6A is a bar chart illustrating expression levels of a1AT in muscle following delivery of a1AT as described in Example 7.

Fig. 6B is a bar chart demonstrating expression levels of epo in muscle following delivery of epo as described in Example 7.

Fig. 6C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to muscle as described in Example 7.

Fig. 6D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of a1AT or epo to muscle as described in Example 7.

15 Detailed Description of the Invention

The present invention provides novel nucleic acid sequences for an adeno-associated virus of serotype 1 (AAV-1). Also provided are fragments of these AAV-1 sequences. Among particularly desirable AAV-1 fragments are the inverted terminal repeat sequences (ITRs), rep and cap. Each of these fragments may be readily utilized, e.g., as a cassette, in a variety of vector systems and host cells. Such fragments may be used alone, in combination with other AAV-1 sequences or fragments, or in combination with elements from other AAV or non-AAV viral sequences. In one particularly desirable embodiment, a cassette may contain the AAV-1 ITRs of the invention flanking a selected transgene. In another desirable embodiment, a cassette may contain the AAV-1 rep and/or cap proteins, e.g., for use in producing recombinant (rAAV) virus.

Thus, the AAV-1 sequences and fragments thereof are useful in production of rAAV, and are also useful as antisense delivery vectors, gene therapy vectors, or vaccine vectors. The invention further provides nucleic acid molecules, gene delivery

10

15

20

25

vectors, and host cells which contain the AAV-1 sequences of the invention. Also provided a novel methods of gene delivery using AAV vectors.

As described herein, the vectors of the invention containing the AAV-1 capsid proteins of the invention are particularly well suited for use in applications in which the neutralizing antibodies diminish the effectiveness of other AAV serotype based vectors, as well as other viral vectors. The rAAV vectors of the invention are particularly advantageous in rAAV readministration and repeat gene therapy.

These and other embodiments and advantages of the invention are described in more detail below. As used throughout this specification and the claims, the term "comprising" is inclusive of other components, elements, integers, steps and the like.

1. AAV-1 NUCLEIC ACID AND PROTEIN SEQUENCES

The AAV-1 nucleic acid sequences of the invention include the DNA sequences of SEQ ID NO: 1 (Figs. 1A-1C), which consists of 4718 nucleotides. The AAV-1 nucleic acid sequences of the invention further encompass the strand which is complementary to SEQ ID NO: 1, as well as the RNA and cDNA sequences corresponding to SEQ ID NO: 1 and its complementary strand. Also included in the nucleic acid sequences of the invention are natural variants and engineered modifications of SEQ ID NO: 1 and its complementary strand. Such modifications include, for example, labels which are known in the art, methylation, and substitution of one or more of the naturally occurring nucleotides with an analog.

Further included in this invention are nucleic acid sequences which are greater than 85%, preferably at least about 90%, more preferably at least about 95%, and most preferably at least about 98 - 99% identical or homologous to SEQ ID NO:1. The term "percent sequence identity" or "identical" in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over the full-length sequence, or a fragment at least about nine nucleotides, usually at least about 20 - 24 nucleotides, at least about 28 - 32 nucleotides, and preferably at least about 36 or more nucleotides. There are a number of different

10

15

20

25

algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using Fasta, a program in GCG Version 6.1. Fasta provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, 1990, herein incorporated by reference). For instance, percent sequence identity between nucleic acid sequences can be determined using Fasta with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference.

The term "substantial homology" or "substantial similarity," when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 95 - 99% of the sequence.

Also included within the invention are fragments of SEQ ID NO: 1, its complementary strand, cDNA and RNA complementary thereto. Suitable fragments are at least 15 nucleotides in length, and encompass functional fragments which are of biological interest. Certain of these fragments may be identified by reference to Figs. 1A-1C. Examples of particularly desirable functional fragments include the AAV-1 inverted terminal repeat (ITR) sequences of the invention. In contrast to the 145 nt ITRs of AAV-2, AAV-3, and AAV-4, the AAV-1 ITRs have been found to consist of only 143 nucleotides, yet advantageously are characterized by the T-shaped hairpin structure which is believed to be responsible for the ability of the AAV-2 ITRs to direct site-specific integration. In addition, AAV-1 is unique among other AAV serotypes, in that the 5' and 3' ITRs are identical. The full-length 5' ITR sequences of AAV-1 are provided at nucleotides l-143 of SEQ ID NO: 1 (Fig. 1A) and the fulllength 3' ITR sequences of AAV-1 are provided at nt 4576-4718 of SEQ ID NO: 1 (Fig. 1C). One of skill in the art can readily utilize less than the full-length 5' and/or 3' ITR sequences for various purposes and may construct modified ITRs using conventional techniques, e.g., as described for AAV-2 ITRs in Samulski et al, Cell,

8

Another desirable functional fragment of the AAV-1 genome is the P5 promoter of AAV-1 which has sequences unique among AAV P5 promoters, while maintaining critical regulatory elements and functions. This promoter is located within nt 236 - 299 of SEQ ID NO: 1 (Fig. 1A). Other examples of functional fragments of interest include the sequences at the junction of the rep/cap, e.g., the sequences spanning nt 2306-2223, as well as larger fragments which encompass this junction which may comprise 50 nucleotides on either side of this junction. Still other examples of functional fragments include the sequences encoding the rep proteins. Rep 78 is located in the region of nt 334 - 2306 of SEQ ID NO: 1; Rep 68 is located in the region of nt 334-2272, and contains an intron spanning nt 1924-2220 of SEQ ID NO: 1. Rep 52 is located in the region of nt 1007 - 2304 of SEQ ID NO: 1; rep 40 is located in the region of nt 1007 - 2272, and contains an intron spanning nt 1924-2246 of SEQ ID NO. 1. Also of interest are the sequences encoding the capsid proteins, VP 1 [nt 2223-4431 of SEQ ID NO: 1], VP2 [nt 2634-4432 of SEQ ID NO: 1] and VP3 [nt 2829-4432 of SEQ ID NO: 1]. Other fragments of interest may include the AAV-1 P19 sequences, AAV-1 P40 sequences, the rep binding site, and the terminal resolute site (TRS).

5

10

15

20

25

The invention further provides the proteins and fragments thereof which are encoded by the AAV-1 nucleic acids of the invention. Particularly desirable proteins include the rep and cap proteins, which are encoded by the nucleotide sequences identified above. These proteins include rep 78 [SEQ ID NO:5], rep 68 [SEQ ID NO:7], rep 52 [SEQ ID NO:9], rep 40 [SEQ ID NO: 11], vpl [SEQ ID NO: 13], vp2 [SEQ ID NO: 15], and vp3 [SEQ IID NO: 17] and functional fragments thereof while the sequences of the rep and cap proteins have been found to be closely related to those of AAV-6, there are differences in the amino acid sequences (see Table 1 below), as well as differences in the recognition of these proteins by the immune system. However, one of skill in the art may readily select other suitable proteins or protein fragments of biological interest. Suitably, such fragments are at least 8 amino acids in length. However, fragments of other desired lengths may be readily utilized.

10

15

20

25

Such fragments may be produced recombinantly or by other suitable means, e.g., chemical synthesis.

The sequences, proteins, and fragments of the invention may be produced by any suitable means, including recombinant production, chemical synthesis, or other synthetic means. Such production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

II. VIRAL VECTORS

In another aspect, the present invention provides vectors which utilize the AAV-1 sequences of the invention, including fragments thereof, for delivery of a heterologous gene or other nucleic acid sequences to a target cell. Suitably, these heterologous sequences (i.e., a transgene) encode a protein or gene product which is capable of being expressed in the target cell. Such a transgene may be constructed in the form of a "minigene". Such a "minigene" includes selected heterologous gene sequences and the other regulatory elements necessary to transcribe the gene and express the gene product in a host cell. Thus, the gene sequences are operatively linked to regulatory components in a manner which permit their transcription. Such components include conventional regulatory elements necessary to drive expression of the transgene in a cell containing the viral vector. The minigene may also contain a selected promoter which is linked to the transgene and located, with other regulatory elements, within the selected viral sequences of the recombinant vector.

Selection of the promoter is a routine matter and is not a limitation of this invention. Useful promoters may be constitutive promoters or regulated (inducible) promoters, which will enable control of the timing and amount of the transgene to be expressed. For example, desirable promoters include the cytomegalovirus (CMV) immediate early promoter/enhancer [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the Rous sarcoma virus LTR promoter/enhancer, and the chicken cytoplasmic β-actin promoter [T. A. Kost et al, Nucl, Acids Res., 11(23):8287 (1983)]. Still other desirable promoters are the albumin promoter and an AAV P5 promoter. Optionally, the selected promoter is used in conjunction with a heterologous enhancer, e.g., the β-

10

actin promoter may be used in conjunction with the CMV enhancer. Yet other suitable or desirable promoters and enhancers may be selected by one of skill in the art.

5

10 .

15

20

25

30

The minigene may also desirably contain nucleic acid sequences heterologous to the viral vector sequences including sequences providing signals required for efficient polyadenylation of the transcript (poly-A or pA) and introns with functional splice donor and acceptor sites. A common poly-A sequence which is employed in the exemplary vectors of this invention is that derived from the papovavirus SV-40. The poly-A sequence generally is inserted in the minigene downstream of the transgene sequences and upstream of the viral vector sequences. A common intron sequence is also derived from SV-40, and is referred to as the SV40 T intron sequence. A minigene of the present invention may also contain such an intron, desirably located between the promoter/enhancer sequence and the transgene. Selection of these and other common vector elements are conventional [see, e.g., Sambrook et al, "Molecular Cloning. A Laboratory Manual", 2d edit., Cold Spring Harbor Laboratory, New York (1989) and references cited therein] and many such sequences are available from commercial and industrial sources as well as from Genebank.

The selection of the transgene is not a limitation of the present invention. Suitable transgenes may be readily selected from among desirable reporter genes, therapeutic genes, and optionally, genes encoding immunogenic polypeptides. Examples of suitable reporter genes include β-galactosidase (β-gal), an alkaline phosphatase gene, and green fluorescent protein (GFP). Examples of therapeutic genes include, cytokines, growth factors, hormones, and differentiation factors, among others. The transgene may be readily selected by one of skill in the art. See, e.g., WO 98/09657, which identifies other suitable transgenes.

Suitably, the vectors of the invention contain, at a minimum, cassettes which consist of fragments of the AAV-1 sequences and proteins. In one embodiment, a vector of the invention comprises a selected transgene, which is flanked by a 5' ITR and a 3' ITR, at least one of which is an AAV-1 ITR of the invention. Suitably,

10

15

20

25

vectors of the invention may contain a AAV-1 P5 promoter of the invention. In yet another embodiment, a plasmid or vector of the invention contains AAV-1 rep sequences. In still another embodiment, a plasmid or vector of the invention contains at least one of the AAV-1 cap proteins of the invention. Most suitably, these AAV-1-derived vectors are assembled into viral vectors, as described herein.

A. AAV Viral Vectors

In one aspect, the present invention provides a recombinant AAV-1 viral vector produced using the AAV-1 capsid proteins of the invention. The packaged rAAV-1 virions of the invention may contain, in addition to a selected minigene, other AAV-1 sequences, or may contain sequences from other AAV serotypes.

Methods of generating rAAV virions are well known and the selection of a suitable method is not a limitation on the present invention. See, e.g., K. Fisher et al., J. Virol., 70:520-532 (1993) and US Patent 5,478,745. In one suitable method, a selected host cell is provided with the AAV sequence encoding a rep protein, the gene encoding the AAV cap protein and with the sequences for packaging and subsequent delivery. Desirably, the method utilizes the sequences encoding the AAV-1 rep and/or cap proteins of the invention.

In one embodiment, the rep/cap genes and the sequences for delivery are supplied by co-transfection of vectors carrying these genes and sequences. In one currently preferred embodiment, a cis (vector) plasmid, a trans plasmid containing the rep and cap genes, and a plasmid containing the adenovirus helper genes are co-transfected into a suitable cell line, e.g., 293. Alternatively, one or more of these functions may be provided in trans via separate vectors, or may be found in a suitably engineered packaging cell line.

An exemplary cis plasmid will contain, in 5' to 3' order, AAV 5' ITR, the selected transgene, and AAV 3' ITR. In one desirable embodiment, at least one of the AAV ITRs is a 143 nt AAV-1 ITR. However, other AAV serotype ITRs may be readily selected. Suitably, the full-length ITRs are utilized. However, one of skill in

. 10

15

20

25

30

the art can readily prepare modified AAV ITRs using conventional techniques.

Similarly, methods for construction of such plasmids is well known to those of skill in the art.

A trans plasmid for use in the production of the rAAV-1 virion particle may be prepared according to known techniques. In one desired embodiment, this plasmid contains the rep and cap proteins of AAV-1, or functional fragments thereof. Alternatively, the rep sequences may be from another selected AAV serotype.

The cis and trans plasmid may then be co-transfected with a wild-type helper virus (e.g., Ad2, Ad5, or a herpesvirus), or more desirably, a replication - defective adenovirus, into a selected host cell. Alternatively, the cis and trans plasmid may be co-transfected into a selected host cell together with a transfected plasmid which provides the necessary helper functions. Selection of a suitable host cell is well within the skill of those in the art and include such mammalian cells as 293 cells, HeLa cells, among others.

Alternatively, the cis plasmid and, optionally the trans plasmid, may be transfected into a packaging cell line which provides the remaining helper functions necessary for production of a rAAV containing the desired AAV-1 sequences of the invention. An example of a suitable packaging cell line, where an AAV-2 capsid is desired, is B-50, which stably expresses AAV-2 rep and cap genes under the control of a homologous P5 promoter. This cell line is characterized by integration into the cellular chromosome of multiple copies (at least 5 copies) of P5-rep-cap gene cassettes in a concatomer form. This B-50 cell line was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, on September 18, 1997 under Accession No. CRL-12401 pursuant to the provisions of the Budapest Treaty. However, the present invention is not limited as to the selection of the packaging cell line.

Exemplary transducing vectors based on AAV-1 capsid proteins have been tested both *in vivo and in vitro*, as described in more detail in Example 4. In these studies, it was demonstrated that recombinant AAV vector with an AAV-1 virion can transduce both mouse liver and muscle. These, and other AAV-1 based

13

gene therapy vectors which may be generated by one of skill in the art are beneficial for gene delivery to selected host cells and gene therapy patients since the neutralization antibodies of AAV-1 present in much of the human population exhibit different patterns from other AAV serotypes and therefore do not neutralize the AAV-1 virions. One of skill in the art may readily prepare other rAAV viral vectors containing the AAV-1 capsid proteins provided herein using a variety of techniques known to those of skill in the art. One may similarly prepare still other rAAV viral vectors containing AAV-1 sequence and AAV capsids of another serotype.

B. Other Viral Vectors

10

5

One of skill in the art will readily understand that the AAV-1 sequences of the invention can be readily adapted for use in these and other viral vector systems for *in vitro*, *ex vivo or in vivo* gene delivery. Particularly well suited for use in such viral vector systems are the AAV-1 ITR sequences, the AAV-1 rep, the AAV-1 cap, and the AAV-1 P5 promoter sequences.

15

20

25

30

For example, in one desirable embodiment, the AAV-1 ITR sequences of the invention may be used in an expression cassette which includes AAV-1 5' ITR, a non-AAV DNA sequences of interest (e.g., a minigene), and 3' ITR and which lacks functional rep/cap. Such a cassette containing an AAV-1 ITR may be located on a plasmid for subsequent transfection into a desired host cell, such as the cis plasmid described above. This expression cassette may further be provided with an AAV capsid of a selected serotype to permit infection of a cell or stably transfected into a desired host cell for packaging of rAAV virions. Such an expression cassette may be readily adapted for use in other viral systems, including adenovirus systems and lentivirus systems. Methods of producing Ad/AAV vectors are well known to those of skill in the art. One desirable method is described in PCT/US95/14018. However, the present invention is not limited to any particular method.

Another aspect of the present invention is the novel AAV-1 P5 promoter sequences which are located in the region spanning nt 236 - 299 of SEQ ID NO: 1. This promoter is useful in a variety of viral vectors for driving expression of a desired transgene.

14

Similarly, one of skill in the art can readily select other fragments of the AAV-1 genome of the invention for use in a variety of vector systems. Such vectors systems may include, e.g., lentiviruses, retroviruses, poxviruses, vaccinia viruses, and adenoviral systems, among others. Selection of these vector systems is not a limitation of the present invention.

C. Host Cells And Packaging Cell Lines

5

10

15

20

25

In yet another aspect, the present invention provides host cells which may be transiently transfected with AAV-1 nucleic acid sequences of the invention to permit expression of a desired transgene or production of a rAAV particle. For example, a selected host cell may be transfected with the AAV-1 P5 promoter sequences and/or the AAV-1 5' ITR sequences using conventional techniques. Providing AAV helper functions to the transfected cell lines of the invention results in packaging of the rAAV as infectious rAAV particles. Such cell lines may be produced in accordance with known techniques [see, e.g, US Patent No. 5,658,785], making use of the AAV-1 sequences of the invention.

Alternatively, host cells of the invention may be stably transfected with a rAAV expression cassette of the invention, and with copies of AAV-1 rep and cap genes. Suitable parental cell lines include mammalian cell lines and it may be desirable to select host cells from among non-simian mammalian cells. Examples of suitable parental cell lines include, without limitation, HeLa [ATCC CCL 2], A549 [ATCC Accession No. CCL 185], KB [CCL 17], Detroit [e.g., Detroit 510, CCL 72] and WI-38 [CCL 75] cells. These cell lines are all available from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 USA. Other suitable parent cell lines may be obtained from other sources and may be used to construct stable cell lines containing the P5 and/or AAV rep and cap sequences of the invention.

Recombinant vectors generated as described above are useful for delivery of the DNA of interest to cells.

15

20

25

III. METHODS OF DELIVERING GENES VIA AAV-1 DERIVED VECTORS

In another aspect, the present invention provides a method for delivery of a transgene to a host which involves transfecting or infecting a selected host cell with a recombinant viral vector generated with the AAV-1 sequences (or functional fragments thereof) of the invention. Methods for delivery are well known to those of skill in the art and are not a limitation of the present invention.

In one desirable embodiment, the invention provides a method for AAV-mediated delivery of a transgene to a host. This method involves transfecting or
infecting a selected host cell with a recombinant viral vector containing a selected
transgene under the control of sequences which direct expression thereof and AAV-1
capsid proteins.

Optionally, a sample from the host may be first assayed for the presence of antibodies to a selected AAV serotype. A variety of assay formats for detecting neutralizing antibodies are well known to those of skill in the art. The selection of such an assay is not a limitation of the present invention. See, e.g., Fisher et al, Nature Med., 3(3):306-312 (March 1997) and W. C. Manning et al, Human Gene Therapy, 9:477-485 (March 1, 1998). The results of this assay may be used to determine which AAV vector containing capsid proteins of a particular serotype are preferred for delivery, e.g., by the absence of neutralizing antibodies specific for that capsid serotype.

In one aspect of this method, the delivery of vector with AAV-1 capsid proteins may precede or follow delivery of a gene via a vector with a different serotype AAV capsid protein. Thus, gene delivery via rAAV vectors may be used for repeat gene delivery to a selected host cell. Desirably, subsequently administered rAAV vectors carry the same transgene as the first rAAV vector, but the subsequently administered vectors contain capsid proteins of serotypes which differ from the first vector. For example, if a first vector has AAV-2 capsid proteins, subsequently administered vectors may have capsid proteins selected from among the other serotypes, including AAV-1, AAV-3A, AAV-3B, AAV-4 and AAV-6.

16

Thus, a rAAV-1-derived recombinant viral vector of the invention provides an efficient gene transfer vehicle which can deliver a selected transgene to a selected host cell *in vivo or ex vivo* even where the organism has neutralizing antibodies to one or more AAV serotypes. These compositions are particularly well suited to gene delivery for therapeutic purposes. However, the compositions of the invention may also be useful in immunization. Further, the compositions of the invention may also be used for production of a desired gene product *in vitro*.

5

10

15

20

25

30

The above-described recombinant vectors may be delivered to host cells according to published methods. An AAV viral vector bearing the selected transgene may be administered to a patient, preferably suspended in a biologically compatible solution or pharmaceutically acceptable delivery vehicle. A suitable vehicle includes sterile saline. Other aqueous and non-aqueous isotonic sterile injection solutions and aqueous and non-aqueous sterile suspensions known to be pharmaceutically acceptable carriers and well known to those of skill in the art may be employed for this purpose.

The viral vectors are administered in sufficient amounts to transfect the cells and to provide sufficient levels of gene transfer and expression to provide a therapeutic benefit without undue adverse effects, or with medically acceptable physiological effects, which can be determined by those skilled in the medical arts. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the liver, oral, intranasal, intravenous, intramuscular, subcutaneous, intradermal, and other parental routes of administration. Routes of administration may be combined, if desired.

Dosages of the viral vector will depend primarily on factors such as the condition being treated, the age, weight and health of the patient, and may thus vary among patients. For example, a therapeutically effective human dosage of the viral vector is generally in the range of from about 1 ml to about 100 ml of solution containing concentrations of from about 1 x 10⁹ to 1 x 10¹⁶ genomes virus vector. A preferred human dosage may be about 1 x 10¹³ to 1 x 10¹⁶ AAV genomes. The dosage will be adjusted to balance the therapeutic benefit against any side effects and

10

15

20

25

such dosages may vary depending upon the therapeutic application for which the recombinant vector is employed. The levels of expression of the transgene can be monitored to determine the frequency of dosage resulting in viral vectors, preferably AAV vectors containing the minigene. Optionally, dosage regimens similar to those described for therapeutic purposes may be utilized for immunization using the compositions of the invention. For *in vitro* production, a desired protein may be obtained from a desired culture following transfection of host cells with a rAAV containing the gene encoding the desired protein and culturing the cell culture under conditions which permits expression. The expressed protein may then be purified and isolated, as desired. Suitable techniques for transfection, cell culturing, purification, and isolation are known to those of skill in the art.

The following examples illustrate several aspects and embodiments of the invention.

Example 1 - Generation of Infectious Clone of AAV-1

The replicated form DNA of AAV-1 was extracted from 293 cells that were infected by AAV-1 and wild type adenovirus type 5.

A. Cell Culture and Virus

AAV-free 293 cells and 84-31 cells were provided by the human application laboratory of the University of Pennsylvania. These cells were cultured in Dulbecco's Modified Eagle Medium with 10% fetal bovine serum (Hyclone), penicillin (100 U/ml) and streptomycin at 37°C in a moisturized environment supplied with 5% CO₂. The 84-31 cell line constitutively expresses adenovirus genes E1a, Elb, E4/ORF6, and has been described previously [K. J. Fisher, J. Virol., 70:520-532 (1996)]. AAV-1 (ATCC VR-645) seed stock was purchased from American Type Culture Collection (ATCC, Manassas, VA). AAV viruses were propagated in 293 cells with wild type Ad5 as a helper virus.

B. Recombinant AAV Generation

The recombinant AAV viruses were generated by transfection using an adenovirus free method. Briefly, the cis plasmid (with AAV ITR), trans plasmid (with

18

AAV rep gene and cap gene) and helper plasmid (pFa13, with essential regions from the adenovirus genome) were simultaneously co-transfected into 293 cells in a ratio of 1:1:2 by calcium phosphate precipitation. The pFa13 helper plasmid has an 8 kb deletion in the adenovirus E2B region and has deletions in most of the late genes. This helper plasmid was generated by deleting the RsrII fragment from pFG140 (Microbix, Canada). Typically, 50 µg of DNA (cis:trans:PFa13 at ratios of 1:1:2, respectively) was transfected onto a 15 cm tissue culture dish. The cells were harvested 96 hours post-transfection, sonicated and treated with 0.5% sodium deoxycholate (37°C for 10 min). Cell lysates were then subjected to two rounds of a CsCl gradient. Peak fractions containing AAV vector were collected, pooled, and dialyzed against PBS before injecting into animals. To make rAAV virus with AAV-1 virion, the pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide rep and cap function.

5

10

15

20

25

30

For the generation of rAAV based on AAV-2, p5E18 was used as the trans plasmid since it greatly improved the rAAV yield. This plasmid, p5E18(2/2), expresses AAV-2 Rep and Cap and contains a P5 promoter relocated to a position 3' to the Cap gene, thereby minimizing expression of Rep78 and Rep68. The strategy was initially described by Li et al, J. Virol., 71:5236-5243 (1997). P5E18(2/2) was constructed in the following way. The previously described pMMTV-trans vector (i.e., the mouse mammary tumor virus promoter substituted for the P5 promoter in an AAV-2-based vector) was digested with Smal and ClaI, filled in with the Klenow enzyme, and then recircularized with DNA ligase. The resulting construct was digested with XbaI, filled in, and ligated to the blunt-ended BamHI-XbaI fragment from pCR-p5, constructed in the following way. The P5 promoter of AAV was amplified by PCR and the amplified fragment was subsequently cloned into pCR2.1 (Invitrogen) to yield pCR-P5. The helper plasmid pAV1H was constructed by cloning the Bfal fragment of pAAV-2 into pBluescript II-SK(+) at the BcorV and Smal sites. The 3.0-kb XbaI-KpnI fragment from p5E18(2/2), the 2.3-kb XbaI-KpnI fragment from pAV1H, and the 1.7-kb KpnI fragment from p5E18(2/2) were incorporated into a separate plasmid P5E18(2/1), which contains AAV-2 Rep, AAV-1 Cap, and the

19

AAV-2 P5 promoter located 3' to the Cap gene. Plasmid p5E18(2/1) produced 10- to 20-fold higher quantities of the vector than pAV1H (i.e., 10¹² genomes/50 15-cm² plates).

C. <u>DNA Techniques</u>

5

10

15

Hirt DNA extraction was performed as described in the art with minor modification [R.J. Samulski et al., Cell, 33:135-143 (1983)]. More particularly, Hirst solution without SDS was used instead of using original Hirt solution containing SDS. The amount of SDS present in the original Hirst solution was added after the cells had been fully suspended. To construct AAV-1 infectious clone, the Hirt DNA from AAV-1 infected 293 cells was repaired with Klenow enzyme (New England Biolabs) to ensure the ends were blunt. The treated AAV-1 Hirt DNA was then digested with BamHI and cloned into three vectors, respectively. The internal BamHI was cloned into pBlueScript II-SK+ cut with BamHI to get pAV1-BM. The left and right fragments were cloned into pBlueScript II-SK+ cut with BamHI + EcoRV to obtain pAV1-BL and pAV1-BR, respectively. The AAV sequence in these three plasmids were subsequently assembled into the same vector to get AAV-1 infectious clone pAAV-1. The helper plasmid for recombinant AAV-1 virus generation was constructed by cloning the Bfa I fragment of pAAV-1 into pBlueScript II-SK+ at the EcoRV site.

20

Analysis of the Hirt DNA revealed three bands, a dimer at 9.4 kb, a monomer at 4.7 kb and single-stranded DNA at 1.7 kb, which correlated to different replication forms of AAV-1. The monomer band was excised from the gel and then digested with *BamH*I. This resulted in three fragments of 1.1 kb, 0.8 kb and 2.8 kb. This pattern is in accordance with the description by Bantel-schaal and zur Hausen, Virol., 134(1):52-63 (1984). The 1.1 kb and 2.8 kb *BamH*I fragments were cloned into pBlueScript-KS(+) at *BamH*I and EcoRV site. The internal 0.8 kb fragment was cloned into *BamH*I site of pBlueScript-KS(+).

These three fragments were then subcloned into the same construct to obtain a plasmid (pAAV-1) that contained the full sequence of AAV-1. The pAAV-1 was then tested for its ability to rescue from the plasmid backbone and package

25

20

infectious virus. The pAAV-1 was then transfected to 293 cells and supplied with adenovirus type as helper at MOI 10. The virus supernatant was used to reinfect 293 cells.

For Southern blot analysis, Hirt DNA was digested with *Dpn*I to remove bacteria-borne plasmid and probed with internal *BamH*I fragment of AAV-1. The membrane was then washed at high stringency conditions, which included: twice 30 minutes with 2X SSC, 0.1% SDS at 65°C and twice 30 minutes with 0.1X SSC, 0.1% SDS at 65°C. The membrane was then analyzed by both phosphor image and X-ray autoradiography. The results confirmed that pAAV-1 is indeed an infectious clone of AAV serotype 1.

Example 2 - Sequencing Analysis of AAV-1

5

10

15

20

25

The entire AAV-1 genome was then determined by automatic sequencing and was found to be 4718 nucleotides in length (Figs. 1A-1C). For sequencing, an ABI 373 automatic sequencer as used to determine the sequences for all plasmids and PCR fragments related to this study using the FS dye chemistry. All sequences were confirmed by sequencing both plus and minus strands. These sequences were also confirmed by sequencing two independent clones of pAV-BM, pAV-BL and pAV-BR. Since the replicated form of AAV-1 DNA served as the template for sequence determination, these sequences were also confirmed by sequencing a series of PCR products using original AAV-1 seed stock as a template.

The length of AAV-1 was found to be within the range of the other serotypes: AAV-3 (4726 nucleotides), AAV-4 (4774 nucleotides), AAV-2 (4681 nucleotides), and AAV-6 (4683 nucleotides).

The AAV-1 genome exhibited similarities to other serotypes of adenoassociated viruses. Overall, it shares more than 80% identity with other known AAV
viruses as determined by the computer program Megalign using default settings
[DNASTAR, Madison, WI]. The key features in AAV-2 can also be found in AAV1. First, AAV-1 has the same type of inverted terminal repeat which is capable of
forming T-shaped hairpin structures, despite the differences at the nucleotide level

10

15

20

25

30

(Figs. 2 and 3). The sequences of right ITRs and left ITRs of AAV-1 are identical. The AAV TR sequence is subdivided into A, A', B, B', C, C', D and D' [Bern, cited above].

These AAV ITR sequences are also virtually the same as those found in AAV-6 right ITR, there being one nucleotide difference in each of A and A' sequence, and the last nucleotide of the D sequence. Second, the AAV-2 rep binding motif [GCTCGCTCGCTCGCTG (SEQ ID NO: 20)] is well conserved. Such motif can also be found in the human chromosome 19 AAV-2 pre-integration region. Finally, non-structural and structural coding regions, and regulatory elements similar to those of other AAV serotypes also exist in AAV-1 genome.

Although the overall features of AAV terminal repeats are very much conserved, the total length of the AAV terminal repeat exhibits divergence. The terminal repeat of AAV-1 consists of 143 nucleotides while those of AAV-2, AAV-3, and AAV-4 are about 145 or 146 nucleotides. The loop region of AAV-1 ITR most closely resembles that of AAV-4 in that it also uses TCT instead of the TTT found in AAV-2 and AAV-3. The possibility of sequencing error was eliminated using restriction enzyme digestion, since these three nucleotides are part of the SacI site (gagete; nt 69-74 of SEQ ID NO: 1). The p5 promoter region of AAV-1 shows more variations in nucleotide sequences with other AAV serotypes. However, it still maintains the critical regulatory elements. The two copies of YY1 [See, Fig. 1A-1C] sites seemed to be preserved in all known AAV serotypes, which have been shown to be involved in regulating AAV gene expression. In AAV-4, there are 56 additional nucleotides inserted between YY1 and E-box/USF site, while in AAV-1, there are 26 additional nucleotides inserted before the E-box/USF site. The p19 promoter, p40 promoter and polyA can also be identified from the AAV-1 genome by analogy to known AAV serotypes, which are also highly conserved.

Thus, the analysis of AAV terminal repeats of various serotypes showed that the A and A' sequence is very much conserved. One of the reasons may be the Rep binding motif (GCTC)₃GCTG [SEQ ID NO: 20]. These sequences appear to be essential for AAV DNA replication and site-specific integration. The same sequence

22

has also been shown to be preserved in a monkey genome [Samulski, personal communication]. The first 8 nucleotides of the D sequence are also identical in all known AAV serotypes. This is in accordance with the observation of the Srivastava group that only the first 10 nucleotides are essential for AAV packaging [X.S. Wang et al, J. Virol., 71:3077-3082 (1997); X.S. Wang et al, J. Virol., 71:1140-1146 (1997)]. The function of the rest of the D sequences still remain unclear. They may be somehow related to their tissue specificities. The variation of nucleotide in B and C sequence may also suggest that the secondary structure of the ITRs is more critical for its biological function, which has been demonstrated in many previous publications.

Example 3 - Comparison of AAV-1 Sequences

5

10

15

20

25

The nucleotide sequences of AAV-1, obtained as described above, were compared with known AAV sequences, including AAV-2, AAV-4 and AAV-6 using DNA Star Megalign. This comparison revealed a stretch of 71 identical nucleotides shared by AAV-1, AAV-2 and AAV-6. See, Figs. 1A-1C.

This comparison further suggested that AAV-6 is a hybrid formed by homologous recombination of AAV-1 and AAV-2. See, Figs. 3A and 3B. These nucleotides divide the AAV-6 genome into two regions. The 5' half of AAV-6 of 522 nucleotides is identical to that of AAV-2 except in 2 positions. The 3' half of AAV-6 including the majority of the rep gene, complete cap gene and 3' ITR is 98% identical to AAV-1.

Biologically, such recombination may enable AAV-1 to acquire the ability to transmit through the human population. It is also interesting to note that the ITRs of AAV-6 comprise one AAV-1 ITR and one AAV-2 ITR. The replication model of defective parvovirus can maintain this special arrangement. Studies on AAV integration have shown that a majority of AAV integrants carries deletions in at least one of the terminal repeats. These deletions have been shown to be able to be repaired through gene conversion using the other intact terminal repeat as a template. Therefore, it would be very difficult to maintain AAV-6 as a homogenous population

23

when an integrated copy of AAV-6 is rescued from host cells with helper virus infection. The AAV-6 with two identical AAV-2 ITRs or two identical AAV-1 ITRs should be the dominant variants. The AAV-6 with two AAV-1 ITRs has been observed by Russell's group [Rutledge, cited above (1998)]. So far there is no report on AAV-6 with two AAV-2 ITRs. Acquirement of AAV-2 P5 promoter by AAV-6 may have explained that AAV-6 have been isolated from human origin while AAV-1 with the same virion has not. The regulation of P5 promoter between different species of AAV may be different *in vivo*. This observation suggests the capsid proteins of AAV were not the only determinants for tissue specificity.

10

15

5

Although it is clear that AAV-6 is a hybrid of AAV-1 and AAV-2, AAV-6 has already exhibited divergence from either AAV-1 or AAV-2. There are two nucleotide differences between AAV-6 and AAV-2 in their first 450 nucleotides. There are about 1% differences between AAV-6 and AAV-1 in nucleotide levels from nucleotides 522 to the 3' end. There also exists a quite divergent region (nucleotide 4486-4593) between AAV-6 and AAV-1 (Figs. 1A-1C). This region does not encode any known proteins for AAVs. These differences in nucleotide sequences may suggest that AAV-6 and AAV-1 have gone through some evolution since the recombination took place. Another possible explanation is that there exists another variant of AAV-1 which has yet to be identified. So far, there is no evidence to rule out either possibility. It is still unknown if other hybrids (AAV-2 to AAV-4, etc.) existed in nature.

20

The coding region of AAV-1 was deduced by comparison with other known AAV serotypes. Table 1 illustrates the coding region differences between AAV-1 and AAV-6. The amino acid residues are deduced according to AAV-2.

25

With reference to the amino acid position of AAV-1, Table 1 lists the amino acids of AAV-1 which have been changed to the corresponding ones of AAV-6. The amino acids of AAV-1 are shown to the left of the arrow. Reference may be made to SEQ ID NO: 5 of the amino acid sequence of AAV-1 Rep 78 and to SEQ ID NO: 13 for the amino acid sequence of AAV-1 VP1.

24

Table 1

Coding region variations between AAV-1 and AAV-6

Rep prote	ein (Rep78)	Cap protein (VP1)	
Position(s)	Amino acids	Position(s)	Amino acids
28	S-N	129	L-F
191	Q→H	418	E-D
192	H-D	531	E→K
308	E-D	584	F-L
		598	A-V
		642	N-H

It was surprising to see that the sequence of the AAV-1 coding region is

almost identical to that of AAV-6 from position 452 to the end of coding region

(99%). The first 508 nucleotides of AAV-6 have been shown to be identical to those

of AAV-2 [Rutledge, cited above (1998)]. Since the components of AAV-6 genome

seemed to be AAV-2 left ITR - AAV-2 p5 promoter - AAV-1 coding region - AAV
1 right ITR, it was concluded that AAV-6 is a naturally occurred hybrid between

AAV-1 and AAV-2.

Example 4 - Gene Therapy Vector Based on AAV-1

Recombinant gene transfer vectors based on AAV-1 viruses were constructed by the methods described in Example 1. To produce a hybrid recombinant virus with AAV-1 virion and AAV-2 ITR, the AAV-1 trans plasmid (pAV1H) and the AAV-2 cis-lacZ plasmid (with AAV-2 ITR) were used. The AAV-2 ITR was used in this vector in view of its known ability to direct site-specific integration. Also constructed for use in this experiment was an AAV-1 vector carrying the green fluorescent protein (GFP) marker gene under the control of the immediate early promoter of CMV using pAV1H as the trans plasmid.

5

20

10

15

20

25

A. rAAV-1 Viruses Transfect Host Cells in Vitro

84-31 cells, which are subclones of 293 cells (which express adenovirus E1a, E1b) which stably express E4/ORF5, were infected with rAAV-1 GFP or rAAV-lacZ. High levels of expression of GFP and lacZ was detected in the cultured 84-31 cells. This suggested that rAAV-1 based vector was very similar to AAV-2 based vectors in ability to infect and expression levels.

B. rAAV-1 Viruses Transfect Cells in Vivo

The performance of AAV-1 based vectors was also tested *in vivo*. The rAAV-1 CMV-α1AT virus was constructed as follows. The EcoRI fragment of pAT85 (ATCC) containing human α1-antitrypsin (α1AT) cDNA fragment was blunted and cloned into PCR (Promega) at a Smal site to obtain PCR-α1AT. The CMV promoter was cloned into PCR-α1AT at the Xbal site. The Alb-α1AT expression cassette was removed by Xhol and ClaI and cloned into pAV1H at the Xbal site. This vector plasmid was used to generate AAV-1-CMV-α1AT virus used in the experiment described below.

For screening human antibodies against AAV, purified AAV virus is lysed with Ripa buffer (10 mM Tris pH 8.2, 1% Triton X-100, 1% SDS, 0.15 M NaCl) and separated in 10% SDS-PAGE gel. The heat inactivated human serum was used at a 1 to 1000 dilution in this assay. The rAAV-1 CMV-α1AT viruses were injected into Rag-1 mice through tail vein injection at different dosages. The concentration of human α1-antitrypsin in mouse serum was measured using ELISA. The coating antibody is rabbit anti-human human α1-antitrypsin (Sigma). The goat-antihuman α1-antitrypsin (Sigma) was used as the primary detection antibodies. The sensitivity of this assay is around 0.3 ng/ml to 30 ng/ml. The expression of human α-antitrypsin in mouse blood can be detected in a very encouraging level. This result is shown in Table 2.

26

Table 2

Human Antitrypsin Expressed in Mouse Liver

Amount of virus injected	Week 2 (ng/ml)	Week 4 (ng/ml)
2x10 ¹⁰ genomes	214.2	171.4
1x10 ¹⁰ genomes	117.8	109.8
5x10 ¹⁰ genomes	64.5	67.8
2.5x10 ¹⁰ genomes	30.9	58.4

rAAV-1 CMV-lacZ viruses were also injected into the muscle of C57BL6 mice and similar results were obtained. Collectively, these results suggested that AAV-1 based vector would be appropriate for both liver and muscle gene delivery.

Example 5 - Neutralizing Antibodies Against AAV-1

Simple and quantitative assays for neutralizing antibodies (NAB) to AAV-1 and AAV-2 were developed with recombinant vectors. A total of 33 rhesus monkeys and 77 normal human subjects were screened.

A. Nonhuman Primates

5

10

15

20

25

Wild-caught juvenile rhesus monkeys were purchased from Covance (Alice, Tex.) and LABS of Virginia (Yemassee, SC) and kept in full quarantine. The monkeys weighed approximately 3 to 4 kg. The nonhuman primates used in the Institute for Human Gene Therapy research program are purposefully bred in the United States from specific-pathogen-free closed colonies. All vendors are US Department of Agriculture class A dealers. The rhesus macaques are therefore not infected with important simian pathogens, including the tuberculosis agent, major simian lentiviruses (simian immunodeficiency virus and simian retroviruses), and cercopithecine herpesvirus. The animals are also free of internal and external parasites. The excellent health status of these premium animals minimized the potential for extraneous variables. For this study, serum was obtained from monkeys prior to initiation of any protocol.

27

NAB titers were analyzed by assessing the ability of serum antibody to inhibit the transduction of reporter virus expressing green fluorescent protein (GFP) (AAV1-GFP or AAV2-GFP) into 84-31 cells. Various dilutions of antibodies preincubated with reporter virus for 1 hour at 37°C were added to 90% confluent cell cultures. Cells were incubated for 48 hours and the expression of green fluorescent protein was measured by FluoroImaging (Molecular Dynamics). NAB titers were calculated as the highest dilution at which 50% of the cells stained green.

Analysis of NAB in rhesus monkeys showed that 61% of animals tested positive for AAV-1; a minority (24%) has NAB to AAV-2. Over one-third of animals had antibodies to AAV-1 but not AAV-2 (i.e., were monospecific for AAV-1), whereas no animals were positive for AAV-2 without reacting to AAV-1. These data support the hypothesis that AAV-1 is endemic in rhesus monkeys. The presence of true AAV-2 infections in this group of nonhuman primates is less clear, since cross-neutralizing activity of an AAV-1 response to AAV-2 can not be ruled out. It is interesting that there is a linear relationship between AAV-2 NAB and AAV-1 NAB in animals that had both.

B. Humans

5

10

15

20

25

For these neutralization antibody assays, human serum samples were incubated at 56°C for 30 min to inactivate complement and then diluted in DMEM. The virus (rAAV or rAd with either lacZ or GFP) was then mixed with each serum dilution (20X, 400X, 2000X, 4000X, etc.) and incubated for 1 hour at 37°C before applied to 90% confluent cultures of 84-31 cells (for AAV) or Hela cells (for adenovirus) in 96-well plates. After 60 minutes of incubation at culture condition, 100 µl additional media containing 20% FCS was added to make final culture media containing 10% FCS.

The result is summarized in Table 3.

Table 3

Adenovirus	AAV-1	AAV-2	# of samples	Percentage
-	-	-	41	53.2%
+	-	-	16	20.8%
-	+	+	0	0.0%
-	_	+	2	2.6%
-	+	+	2	2.6%
+ .′	-	+	3	3.9%
+ .	. +	-	0 ,	0.0%
+	+	+	13	16.9%
		Total	77	100%

10

15

20

25

5

The human neutralizing antibodies against these three viruses seemed to be unrelated since the existence of neutralizing antibodies against AAV are not indications for antibodies against adenovirus. However, AAV requires adenovirus as helper virus, in most of the cases, the neutralizing antibodies against AAV correlated with the existence of neutralizing antibodies to adenovirus. Among the 77 human serum samples screened, 41% of the samples can neutralize the infectivity of recombinant adenovirus based on Ad5. 15/77 (19%) of serum samples can neutralize the transduction of rAAV-1 while 20/77 (20%) of the samples inhibit rAAV-2 transduction at 1 to 80 dilutions or higher. All serum samples positive in neutralizing antibodies for AAV-1 in are also positive for AAV-2. However, there are five (6%) rAAV-2 positive samples that failed to neutralize rAAV-1. In samples that are positive for neutralizing antibodies, the titer of antibodies also varied in the positive ones. The results from screening human sera for antibodies against AAVs supported the conclusion that AAV-1 presents the same epitome as that of AAV-2 to interact

10

15

20

25

with cellular receptors since AAV-1 neutralizing human serums can also decrease the infectivity of AAV-2. However, the profile of neutralizing antibodies for these AAVs is not identical, there are additional specific receptors for each AAV serotype.

Example 6 - Recombinant AAV Viruses Exhibit Tissue Tropism

The recombinant AAV-1 vectors of the invention and the recombinant AAV-2 vectors [containing the gene encoding human α 1-antitrypsin (α 1AT) or murine erythropoietin (Epo) from a cytomegalovirus-enhanced β -actin promoter (CB)] were evaluated in a direct comparison to equivalent copies of AAV-2 vectors containing the same vector genes.

Recombinant viruses with AAV-1 capsids were constructed using the techniques in Example 1. To make rAAV with AAV-1 virions, pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide Rep and Cap functions. For the generation of the rAAV based on AAV-2, p5E18(2/2) was used as the *trans* plasmid, since it greatly improved the rAAV yield. [Early experiments indicated similar *in vivo* performances of AAV-1 vectors produced with pAV1H and p5E19 (2/1). All subsequent studies used AAV-1 vectors derived from p5E18(2/1) because of the increased yield.]

Equivalent stocks of the AAV-1 and AAV-2 vectors were injected intramuscularly (5 x 10¹⁰ genomes) or liver via the portal circulation (1 x 10¹¹ genomes) into immunodeficient mice, and the animals (four groups) were analyzed on day 30 for expression of transgene. See, Figs. 4A and 4B.

AAV-2 vectors consistently produced 10- to 50-fold more serum erythropoietin or a1-antitrypsin when injected into liver compared to muscle. (However, the AAV-1-delivered genes did achieve acceptable expression levels in the liver.) This result was very different from that for AAV-1 vectors, with which muscle expression was equivalent to or greater than liver expression. In fact, AAV-1 outperformed AAV-2 in muscle when equivalent titers based on genomes were administered.

30

Example 7 - Gene Delivery via rAAV-1

5

10

15

20

25

30

C57BL/6 mice (6- to 8-week old males, Jackson Laboratories) were analyzed for AAV mediated gene transfer to liver following intrasplenic injection of vector (i.e., targeted to liver). A total of 10¹¹ genome equivalents of rAAV-1 or rAAV-2 vector were injected into the circulation in 100 μl buffered saline. The first vector contained either an AAV-1 capsid or an AAV-2 capsid and expressed α1AT under the control of the chicken β-actin (CB) promoter. Day 28 sera were analyzed for antibodies against AAV-1 or AAV-2 and serum α1AT levels were checked. Animals were then injected with an AAV-1 or AAV-2 construct expressing erythropoietin (Epo, also under the control of the CB promoter). One month later sera was analyzed for serum levels of Epo. The following groups were analyzed (Figs. 5A-5D).

In Group 1, vector 1 was AAV-2 expressing a1AT and vector 2 was AAV-2 expressing Epo. Animals generated antibodies against AAV-2 following the first vector administration which prevented the readministration of the AAV-2 based vector. There was no evidence for cross-neutralizing the antibody to AAV-1.

In Group 2, vector 1 was AAV-1 expressing a1AT while vector 2 was AAV-1 expressing Epo. The first vector administration did result in significant a1AT expression at one month associated with antibodies to neutralizing antibodies to AAV-1. The animals were not successfully readministered with the AAV-1 Epo expressing construct.

In Group 3, the effectiveness of an AAV-2 vector expressing Epo injected into a naive animal was measured. The animals were injected with PBS and injected with AAV-2 Epo vector at day 28 and analyzed for Epo expression one month later. The neutralizing antibodies were evaluated at day 28 so we did not expect to see anything since they received PBS with the first vector injection. This shows that in naive animals AAV-2 is very efficient at transferring the Epo gene as demonstrated by high level of serum Epo one month later.

Group 4 was an experiment similar to Group 3 in which the animals originally received PBS for vector 1 and then the AAV-1 expressing Epo construct 28 days later. At the time of vector injection, there obviously were no antibodies to either

10

15

20

AAV-1 or AAV-2. The AAV-1 based vector was capable of generating significant expression of Epo when measured one month later.

Group 5 is a cross-over experiment where the initial vector is AAV-2 expressing α1AT followed by the AAV-1 construct expressing Epo. The animals, as expected, were efficiently infected with the AAV-2 vector expressing α1AT as shown by high levels of the protein in blood at 28 days. This was associated with significant neutralizing antibodies to AAV-2. Importantly, the animals were successfully administered AAV-1 following the AAV-2 vector as shown by the presence of Epo in serum 28 days following the second vector administration. At the time of this vector administration, there was high level AAV-2 neutralizing antibodies and very low cross-reaction to AAV-1. The level of Epo was slightly diminished possibly due to a small amount of cross-reactivity. Group 6 was the opposite cross-over experiment in which the initial vector was AAV-1 based, whereas the second experiment was AAV-2 based. The AAV-1 vector did lead to significant gene expression of α1AT, which also resulted in high level AAV-1 neutralizing antibody. The animals were very efficiently administered AAV-2 following the initial AAV-1 vector as evidenced by high level Epo.

A substantially identical experiment was performed in muscle in which 5 x 10¹⁰ genomes were injected into the tibialis anterior of C57BL/6 mice as a model for muscle directed gene therapy. The results are illustrated in Figs. 6A-6D and are essentially the same as for liver.

In summary, this experiment demonstrates the utility of using an AAV-1 vector in patients who have pre-existing antibodies to AAV-2 or who had initially received an AAV-2 vector and need readministration.

25 Example 8 - Construction of Recombinant Viruses Containing AAV-1 ITRs

This example illustrates the construction of recombinant AAV vectors which contain AAV-1 ITRs of the invention.

An AAV-1 cis plasmid is constructed as follows. A 160 bp Xho-NruI AAV-1 fragment containing the AAV-1 5' ITR is obtained from pAV1-BL pAV1-BL was

generated as described in Example 1. The Xho-NruI fragment is then cloned into a second pAV1-BL plasmid at an XbaI site to provide the plasmid with two AAV-1 ITRs. The desired transgene is then cloned into the modified pAV-1BL at the NruI and BamHI site, which is located between the AAV-1 ITR sequences. The resulting AAV-1 cis plasmid contains AAV-1 ITRs flanking the transgene and lacks functional AAV-1 rep and cap.

5

10

15

20

25

Recombinant AAV is produced by simultaneously transfecting three plasmids into 293 cells. These include the AAV-1 cis plasmid described above; a trans plasmid which provides AAV rep/cap functions and lacks AAV ITRs; and a plasmid providing adenovirus helper functions. The rep and/or cap functions may be provided in trans by AAV-1 or another AAV serotype, depending on the immunity profile of the intended recipient. Alternatively, the rep or cap functions may be provided in cis by AAV-1 or another serotype, again depending on the patient's immunity profile.

In a typical cotransfection, 50 µg of DNA (cis:trans:helper at ratios of 1:1:2, respectively) is transfected onto a 15 cm tissue culture dish. Cells are harvested 96 hours post transfection, sonicated and treated with 0.5% sodium deoxycholate (37° for 10 min). Cell lysates are then subjected to 2-3 rounds of ultracentrifugation in a cesium gradient. Peak fractions containing rAAV are collected, pooled and dialyzed against PBS. A typical yield is 1 x 10¹³ genomes/10⁹ cells.

Using this method, one recombinant virus construct is prepared which contains the AAV-1 ITRs flanking the transgene, with an AAV-1 capsid. Another recombinant virus construct is prepared with contains the AAV-1 ITRs flanking the transgene, with an AAV-2 capsid.

All publications cited in this specification are incorporated herein by reference. While the invention has been described with reference to a particularly preferred embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the claims.

What is claimed is:

- 1. An isolated AAV-1 nucleic acid molecule comprising a sequence selected from the group consisting of:
 - (a) SEQ ID NO: 1;
 - (b) a DNA sequence complementary to SEQ ID NO: 1;
 - (c) cDNA complementary to (a) or (b); and
 - (d) RNA complementary to any of (a) to (c).
- 2. A nucleic acid molecule comprising an AAV-1 inverted terminal repeat (ITR) sequence selected from the group consisting of:
 - (a) nt 1 to 143 of SEQ ID NO: 1;
 - (b) nt 4576 to 4718 of SEQ ID NO: 1;
 - (c) a nucleic acid sequence complementary to (a) or (b); and
 - (d) a functional fragment of (a), (b), or (c).
- 3. A recombinant vector comprising a 5' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:
 - (a) nt 1 to 143 of SEQ ID NO: 1;
 - (b) a nucleic acid sequence complementary to (a), and
 - (c) a functional fragment of (a) or (b).
- 4. The recombinant vector according to claim 3, wherein said vector further comprises a 3' AAV-1 ITR.

34

5. A recombinant vector comprising a 3' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:

- (a) nt 4576 to 4718 of SEQ ID NO: 1;
- (b) a nucleic acid sequence complementary to (a); and
- (c) a functional fragment of (a) or (b).
- 6. The recombinant vector according to claim 5, wherein said vector further comprises a 5' AAV-1 ITR.
- 7. The recombinant vector according to any of claims 3-6, wherein said vector further comprises AAV-1 capsid proteins having the sequence of SEQ ID NO: 13, 15 or 17 or functional fragments thereof.
- 8. The recombinant vector according to any of claims 3-6, wherein said vector further comprises adenovirus sequences.
- 9. A recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.
- 10. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said cap coding region comprises at least one member is selected from the group consisting of:
 - (a) vpl, nt 2223 to 4431 of SEQ ID NO: 1;
 - (b) vp2, nt 2634 to 4432 of SEQ ID NO: 1; and
 - (c) vp3, nt 2829 to 4432 of SEQ ID NO: 1.

35

- 11. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said rep coding region comprises an AAV-1 rep coding region comprising at least one member selected from the group consisting of:
 - (a) rep 78, nt 335 to 2304 of SEQ ID NO: 1;
- (b) rep 68, nt 335 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto;
 - (c) rep 52, nt 1007 to 2304 of SEQ ID NO: 1; and
- (d) rep 40, nt 1007 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto.
- 12. A host cell transduced with a recombinant viral vector according to any of claims 3-6.
- 13. A host cell transduced with a nucleic acid molecule according to any of claims 1, 2, 10 or 11.
- 14. A host cell stably transduced with an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1.
- 15. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to any of claims 3-6.
- 16. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to claim 7.
- 17. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to claim 8.

- 18. A method for AAV-mediated delivery of a transgene comprising the step of delivering to a host cell an AAV virion which comprises:
- (a) a capsid comprising at least one capsid protein encoded by an AAV-1 cap gene; and
- (b) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 19. A method for AAV-mediated delivery of a transgene to a host comprising the steps of:
- (a) assaying a sample from the host to determine the presence of neutralizing antibodies specific against any serotype of AAV; and
 - (b) delivering to the host an AAV virion which comprises:
- (i) a capsid comprising at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has no antibodies as determined in step (a); and
- (ii) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 20. The method according to claim 19, comprising the additional step of repeating steps (a) and (b).
- 21. Use of an AAV virion which comprises a capsid comprising (a) at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has antibodies, and (b) a DNA molecule comprising a transgene operably linked to regulatory sequences directing its expression,

in the preparation of a medicament for delivery of a transgene to a host, wherein said host has no preexisting neutralizing antibodies against the AAV serotype of said cap gene.

37

22. A method for delivery of a transgene comprising the step of delivering to a host cell a recombinant virus comprising a recombinant vector according to any of claims 3-8.

23. A method for producing a selected gene product comprising the steps of transfecting a mammalian cell with the molecule according to claim 1 or a functional fragment thereof and culturing said cell under conditions suitable to express said gene product.

FIG 1A

AAV-2 .	tgcccactccctctctgcgcgctcgctcgctcggtggggcctgcggaccaaaggtccgcgg.gcgcgacag.gcgc.	60 60 60
AAV-2 c	gacggcagagetetgetetgetgetgetgetgetgetgetgetgetget	120 120 120
TR	S `	
AAV-1 g	gcaactccatcactaggggtaaTCGCGAAGCGCCTCCCACGCTGCCGCGTCAGCGCTGA	180
カカなた。つ	Ct .GGTG .AG	163
AAV-6 .	cTG.AG	163
	E box/USF	
N N 17 . 7 . C	GTAAATTACGTCATAGGGGAGTGGTCCTGTATTAGCTGTCACGTGAGTGCTTTTGC	237
AAV-1 C	GAGAG	222
AAV-2 .	G	222
AAV-O.		
	YY1 P <u>5/TATA</u>	
AAV-1 G	ACATTTTGCGACACCACGTGGCCATTTAGGGTATATATGGCCGAGTGAGCGAGC	297
AAV-2	TT.CGCTT.A.CACAC	282
AAV-6.	TTCGCTTA.CACG.	282
7	Y1/p5 RNA Rep_78/68	356
AAV-1 C	TCCATTTTGAC-CGCGAAATTTGAACGAGCAGCAGCCATGCCGGGCTTCTACGAGATCG	342
AAV-2		341
AAV-6		241
N N S7 1 - F	rgatcaaggtgccgagcgacctggacgagcacctgccgggcatttctgactcgtttgtga	416
MMV-1	TCCTGTCAGC	402
AAV-6	TCC	401
		47
AAV-1	GCTGGGTGGCCGAGAAGGAATGGGAGCTGCCCCCGGATTCTGACATGGATCTGAATCTGA	476
AAV-2	A	461
AAV-6	A	40.
1 – זותה	TTGAGCAGGCACCCTGACCGTGGCCGAGAAGCTGCAGCGCGACTTCCTGGTCCAATGGC	536
DAV-2	TACGG	522
AAV-6		52:
AAV-1	GCCGCGTGAGTAAGGCCCCGGAGGCCCTCTTCTTTGTTCAGTTCGAGAAGGGCGAGTCCT	59
AAV-2	TAAG	583
AAV-6	••••••	58
	ACTTCCACCTCCATATTCTGGTGGAGACCACGGGGGTCAAATCCATGGTGCTGGGCCGCT	65
AAV-1	A.GCG.GCACGTTAT.	64
AAV-Z		64
AAV-0		
AAV-1	TCCTGAGTCAGATTAGGGACAAGCTGGTGCAGACCATCTACCGCGGGATCGAGCCGACCC	71
AAV-2	TTTT	70
AAV-6		70
		77
AAV-1	TGCCCAACTGGTTCGCGGTGACCAAGACGCGTAATGGCGCCGGAGGGGGAACAAGGTGG	76
AAV-2	A	76

FIG 1B

2 – עממ	TGGACGAGTGCTACATCCCCAACTACCTCCTGCCCAAGACTCAGCCCGAGCTGCAGTGGGTT.GCACTC	836 822 821
7 7 T T T T	P19/TATA P19 RNA CGTGGACTAACATGGAGGAGTATATAAGCGCCTGTTTGAACCTGGCCGAGCGCAAACGGCTACTCGTCA.GTTCGGAGACGG.C	896 882 881
7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	TCGTGGCGCAGCACCTGACCCACGTCAGCCAGACCCAGGAGCAGAACAAGGAGAATCTGA .GTGGTCGGAA	956 942 941
7777-2	Ψ	1016 1002 1001
7 7 TT 2	TCGGGTGGCTGGTGGACCGGGGCATCACCTCCGACIMGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGT	1076 1062 1061
7 7 T T 7	CCTCGTACATCTCCTTCAACGCCGCTTCCAACTCGCGGTCCCAGATCAAGGCCGCTCTGGATGCATCT	1136 1122 1121
2 – 17 ת ת	ACAATGCCGGCAAGATCATGGCGCTGACCAAATCCGCGCCCGACTACCTGGTAGGCCCCG	1196 1182 1181
AAV-1	CTCCGCCCGCGGACATTAAAACCAACCGCATCTACCGCATCCTGGAGCTGAACGGCTACG AGCGTG.ATCC.GTGTTAAATTAAG	1256 1242 1241
AAV-6	AACCTGCCTACGCCGGCTCCGTCTTTCTCGGCTGGGCCCAGAAAAGGTTCGGGAAGCGCA TCCAATG.CTGAACAAA	1301
	1 ACACCATCTGGCTGTTTGGGCCGGCCACCACGGGCAAGACCAACATCGCGGAAGCCATCG 2	1002
AAV-	1 CCCACGCCGTGCCCTTCTACGGCTGCGTCAACTGGACCAATGAGAACTTTCCCTTCAATG 2A.T	142
AAV-	1 ATTGCGTCGACAAGATGGTGATCTGGTGGGAGGGGGGAAGATGACGGCCAAGGTCGTGG 2 .CT	148
AAV- AAV-	1 AGTCCGCCAAGGCCATTCTCGGCGGCAGCAAGGTGCGCGTGGACCAAAAGTGCAAGTCGT 2GAAA	154
AAV- AAV-	1 CCGCCCAGATCGACCCCACCCCGTGATCGTCACCTCCAACACCAACATGTGCGCCGTGA	161 160

FIG 1C

AAV-1	TTGACGGGAACAGCACCACCTTCGAGCACCAGCAGCCGTTGCAGGACCGGATGTTCAAAT	16/6
AAV-2		1662 1661
AAV-6		TOOT
ת <i>ת</i> ז ז ת ת	TTGAACTCACCCGCCGTCTGGAGCATGACTTTGGCAAGGTGACAAAGCAGGAAGTCAAAG	1740
WWA-1	TIGAACICACCCGCCGICIGGAGCAIGACIIICCGILICCIICCIIICCI	1722
77V-6		1721
AAV-0		
I-VAA	AGTTCTTCCGCTGGGCGCAGGATCACGTGACCGAGGTGGCGCATGAGTTCTACGTCAGAA	1796
AAV-2	.CTGAAAA	1782
AAV-6	***************************************	1781
	P40/TATA	
AAV-1	AGGGTGGAGCCAACAAAAGACCCGCCCCGATGACGCGGATAAAAGCCGAGCCCAAGCGGG	1856
AAV-2		1842
AAV-6		1841
	P40 RNA	1016
AAV-1	CCIGCCCICAGICGCGGAICGAICGAIGGIGGIGGAIGGA	1916 1899
AAV-2	10CGAG	1901
AAV-6		1901
	CCGACAGGTACCAAAACAAATGTTCTCGTCACGCGGGCATGCTTCAGATGCTGTTTCCCT	1976
AAV-1	.A	1959
AAV-2	.п.	1961
AAV-6		
AAV-1	GCAAGACATGCGAGAGAATGAATCAGAATTTCAACATTTGCTTCACGCACG	2036
AAV-2	GACA	2019
AAV-6	<u>~</u>	2021
AAV-1	ACTGTTCAGAGTGCTTCCCCGGCGTGTCAGAATCTCAACCGGTCGTCAGAAAGAGGA	2093
AAV-2	TT	2076
AAV-6	AT	2078
AAV-1	CGTATCGGAAACTCTGTGCCATTCATCATCTGCTGGGGCGGGC	2153
AAV-2	AGCTAA.CAAAATGACA	2133
AAV-6		2138
	Day 70 stan	
_	Rep 78 stop	2213
AAV-1	CGGCCTGCGATCTGGTCAACGTGGACCTGGATGACTGTGTTTCTGAGCAATAAATGACTT	2103
AAV-2	T	2193
AAV-6)	
	∇ VP1 ∇ Rep <u>68</u> st	top
1 <i>– זו</i> ת ת	AAACCAGGTATGGCTGCCGATGGTTATCTTCCAGATTGGCTCGAGGACAACCTCTCTGAG	
מבער בייאמ	T	2253
DAV-6		2258
AAV C	· · · · · · · · · · · · · · · · · · ·	
1 – עום	GGCATTCGCGAGTGGTGGGACTTGAAACCTGGAGCCCGAAGCCCAAAGCCAACCAGCAA	2333
. ₩₩ 1	A. AA.ACA.GC.CCC.AACCAAGCGC	2313
77.V-2		2318
מאת בי	,	
1 – 17 מ מ	AAGCAGGACGACGGCCGGGGTCTGGTGCTTCCTGGCTACAAGTACCTCGGACCCTTCAAC	2393
77.0-3	C.TAAAT	2373
DAW-6	5	2378
4444	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	

FIG 1D

AAV-2	GGACTCGACAAGGGGGAGCCCGTCAACGCGGCGGACGAGCAGGGCCCTCGAGCACAAG	2433
AAV-6	T	2438
AAV-1	GCCTACGACCAGCAGCTCAAAGCGGGTGACAATCCGTACCTGCGGTATAACCACGCCGACGG.CAGCACCAAC	2513 2493
AAV-6		2498
AAV-1	GCCGAGTTTCAGGAGCGTCTGCAAGAAGATACGTCTTTTGGGGGCAACCTCGGGCGAGCAGA	2573 2553
AAV-2 AAV-6	C	_
AAV-1	GTCTTCCAGGCCAAGAAGCGGGTTCTCGAACCTCTCGGTCTGGTTGAGGAAGGCGCTAAG	2633 2613
AAV-6		2618
AAV-1	VP2 ACGGCTCCTGGAAAGAACGTCCGGTAGAGCAGTCGCCACAAGAGCCAGACTCCTCCTCG	2693
AAV-2	GA.GA.G	2673 2678
AAV-1	GGCATCGGCAAGACAGGCCAGCAGCCCGCTAAAAAGAGACTCAATTTTGGTCAGACTGGCA.CAG.GTA.GAT.GA	2753 2733
AAV-6	T	2738
		2813 2793
AAV-2 AAV-6	GCATCCGC.GAGTG. TGCCCAAG.ATAG	2798
	VP3	2873
AAV-2	GTGGGACCTACTACAATGGCTTCAGGCGGTGGCGCACCAATGGCAGACAATAACGAAGGC CAAGAAA	2853
AAV-6		2858
		2933 2913
AAV-2	AA	
DDV-2	AGAGTCATCACCACCAGCACCCGCACCTGGGCCTTGCCCACCTACAATAACCACCTCTAC	2973
AAV-6	TC	2978
AAV-1	AAGCAAATCTCCAGTGCTTCAACGGGGGCCAGCAACGACAACCACTACTTCGGCTACAGC	3053
AAV-2	P A T CCAA A TCG T	3030
	5	
AAV-1	ACCCCTGGGGGTATTTTGATTTCAACAGATTCCACTGCCACTTTTCACCACGTGACTGG	3113
AAV-2	2T	3098
1–עממ	CAGCGACTCATCAACAACTTGGGGATTCCGGCCCAAGAGACTCAACTTCAAACTCTTC	3173
AAV-2	P. AA	3150
AAV-	5	3158
AAV-1	AACATCCAAGTCAAGGAGGTCACGACGAATGATGGCGTCACAACCATCGCTAATAACCTT	3233
AAV-2	2TA	3210

FIG 1E

AAV-1	ACCAGCACGGTTCAAGTCTTCTCGGACTCGGAGTACCAGCTTCCGTACGTCCTCGGCTCT	3293 3270
AAV-2		3278
AAV-6		
1 – 17 ת ת	GCGCACCAGGGCTGCCTCCCTCCGTTCCCGGCGGACGTGTTCATGATTCCGCAATACGGC	3353
AAV-1 AAV-2	TAAGAACG.GAG.TA	3330
MMV-2	G	3338
AAV-0		
1~V44	TACCTGACGCTCAACAATGGCAGCCAAGCCGTGGGACGTTCATCCTTTTACTGCCTGGAA	3413
AAV-2	CCG	3390
AAV-6	A	3398
	·	
AAV-1	TATTTCCCTTCTCAGATGCTGAGAACGGGCAACAACTTTACCTTCAGCTACACCTTTGAG	3473
2-V44	C. T	3450
AAV-6	AG	3458
AAV-1	GAAGTGCCTTTCCACAGCAGCTACGCCCACAGCCAGAGCCTGGACCGGCTGATGAATCCT	3533
AAV-2		3510
AAV-6	C	3498
		2502
AAV-1	CTCATCGACCAATACCTGTATTACCTGAACAGAACTCAAAATCAGTCCGGAAGTGCCCAA	3570
AAV-2	The state of the s	3578
AAV-6		3376
	AACAAGGACTTGCTGTTTAGCCGTGGGTCTCCAGCTGGCATGTCTGTTCAGCCCAAAAAC	3653
AAV-1		3630
AAV-2	(. ()) . AA() (3638
AAV-6		
2211-1	TGGCTACCTGGACCCTGTTATCGGCAGCAGCGCGTTTCTAAAACAAAAACAGACAACAAC	3713
AAV-2		3690
AAV-6		3698
AAV-1	AACAGCAATTTTACCTGGACTGGTGCTTCAAAATATAACCTCAATGGGCGTGAATCCATC	3773
2-1/44	TG A ACT GA.A.C.,GCCCA.ACTC.G	3/50
AAV-6	TA	3758
AAV-1	ATCAACCCTGGCACTGCTATGGCCTCACACAAAGACGACGAAGACAAGTTCTTTCCCATG	3833
AAV-2	G.GTGGC.CCAAGCGTATT	3810
AAV-6		3818
	·	
AAV-1	AGCGGTGTCATGATTTTTGGAAAAGAGAGCGCCGGAGCTTCAAACACTGCATTGGACAAT	3893
DDV-2	G. TC.C. C. G. GC.AG. T.A.AGAAAATGTGAACA.TAG	3870
AAV-6		3878
		2051
AAV-1	GTCATGATTACAGACGAAGAGGAAATTAAAGCCACTAACCCTGTGGCCACCGAAAGATTT	3933
AAV-2		3930
AAV-6		3936
		4013
AAV-1	GGGACCGTGGCAGTCAATTTCCAGAGCAGCAGCAGCAGACCCTGCGACCGGAGATGTGCAT	3001
AAV-2	TT.TAT.TACCCAGAG.C.AG.ATCC	3998
AAV-6	5T) د ر د
n n 7 7 7	GCTATGGGAGCATTACCTGGCATGGTGTGGCAAGATAGAGACGTGTACCTGCAGGGTCCC	4073
AAV-1	A.ACAAC.TTC.TACGCTTG	4050
AAV-2	T	4058
44V-7	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	

FIG 1F

AAV-2	ATTTGGGCCAAAATTCCTCACACAGATGGACACTTTCACCCGTCTCCTCTTATGGGCGGCCAGAGCTACC	4133 4110 4118
AAV-2	.,	4193 4170 4178
AAV-2	CCTCCGGCGGAGTTTTCAGCTACAAAGTTTGCTTCATTCA	4253 4230 4238
AAV-2	CA-AGTGAGTGTGGAAATTGAATGGGAGCTGCAGAAAGAAAACAGCAAGCGCTGGAATCCCGCGCG	4290
AAV-2	CGAAGTGCAGTACACATCCAATTATGCAAAATCTGCCAA-CGTTGATTTTACTGTGGACAA.TTCCAACGTTTGCTTTC	4350
2-עממ	ACAATGGACTTTATACTGAGCCTCGCCCCATTGGCACCCGTTACCTTACCCGTCCCCTGT CTCG.GT.AA.AGTAAT	4431 4410 4416
AAV-1	VP1-3 stop PolyA signal AATTACGTGTTAATCAATAAACCGGTTGATTCGTTTCAGTTGAACTTTGGTCTCCTGTCCG.TT.ATGCGTAGTAGAG	44/0
2-1744	TTCTTATCTTATC-GGTTACCATGGTTAT-AGCTTACACATTAACTGCTTGGTTGCGC .TC.TTATCCGTAGAAGT.GC.TGG.G.GAA.CATTA .ATCA.CA.C-C.GAA	4530
7777_7	TTCGCGATAAAAGACTTACGTCATCGGGttacccctagtgatggagttgcccactccctc ACTA.A.gg.a	4570
AAV-2	tctgcgcgctcgctcggtggggccggcagagcagagctctgccgtctgcggacctt.cacacagccaggca.gc.c.gg	4630
AAV-2	tggtccgcaggccccaccgagcgagcgagcgcgcagagagggagtgggcaacc.g.gctgt	4718 4681 4683

---AAV-1 ITR --aav-1 C G

QC CQ

C G

AA TQ A CA

A CGGGGTGGCTCGCTCGCGCGTCTCCCTCACCCGTT

AAV-1 TR








FIG. 3A




F16. 3B










Group	1	2	3	4	5	6
Vector1 - al AT	AAV2	AAV1	PBS	PBS	AAV2	AAV1
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2



Group	1	2	3	4	5	6
Vector1 - α1AT	AAV2	AAV1	PBS	PBS	AAV2	AAV1
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2

SEQUENCE LISTING

<110> Wilson, James M.

Xiao, Weidong

The Trustees of the University of Pennsylvania

<120> Adeno-Associated Virus Serotype I Nucleic Acid Sequences, Vectors and Host Cells Containing Same

<130> GNVPN.031PCT

<140>

<141>

<150> 60/107,114

<151> 1998-11-05

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 4718

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (335)..(2206)

<220>

<221> CDS

<222> (2223)..(4430)

<400> 1

agacggcaga gctctgctct gccggcccca ccgagcgagc gagcgcgcag agagggagtg 120
ggcaactcca tcactagggg taatcgcgaa gcgcctccca cgctgccgcg tcagcgctga 180
cgtaaattac gtcatagggg agtggtcctg tattagctgt cacgtgagtg cttttgcgac 240
attttgcgac accacgtggc catttagggt atatatggcc gagtgagcga gcaggatctc 300
cattttgacc gcgaaatttg aacgagcagc agcc atg ccg ggc ttc tac gag atc 355
Met Pro Gly Phe Tyr Glu Ile

1

-																403
		_												att Ile		403
Val	TTE	10	vaı	PIO	361	Yab	15	ASP	GIU	1113	ысц	20	Gry	110		
gac	tcg	ttt	gtg	agc	tgg	gtg	gcc	gag	aag	gaa	tgg	gag	ctg	ccc	ccg	451
Asp	Ser	Phe	Val	Ser	Trp	Val	Ala	Glu	Lys	Glu	Trp	Glu	Leu	Pro	Pro	
	25					30					35				-	
4		~		~~+	at a	-	cta	200	~ a ~	636	ac 2	666	cta	acc	ata	499
-		_	_	_	-									Thr		400
40		, iop		,.op	45	7.0				50		•			55	
_		-	-											gtg		547
Ala	Glu	Lys	Leu		Arg	Asp	Phe	Leu		Gln	Trp	Arg	Arg	Val	Ser	
				60				•	65					70		
	~~~		~~~	~~~	at a	***	+++	ätt	C 2 C	ttc	a 2 a	220	aac	gag	tcc	595
_	-	-		-										Glu		373
гуэ	AIa	FIO	75	YIG	Dea	riic	The	80	0111	1110	014	ny o	85	014	001	
					,											
tac	ttc	cac	ctc	cat	att	ctg	gtg	gag	acc	acg	ggg	gtc	aaa	tcc	atg	643
Tyr	Phe	His	Leu	His	Ile	Leu	Val	Glu	Thr	Thr	Gly	Val	Lys	Ser	Met	
		90					95					100				
																601
	_		_											cag		691
Val	Leu 105	GLY	Arg	Pne	ren	110	GIN	116	Arg	Asp	115	Leu	vaı	Gln	1111	
	103					110					110					,
atc	tac	cgc	ggg	atc	gag	ccg	acc	ctg	ccc	aac	tgg	ttc	gcg	gtg	acc	739
														Val		
120					125					130					135	
-	_	-												gag		787
Lys	Thr	Arg	Asn		Ala	Gly	GIŸ	GIŸ	Asn 145	гуs	vai	vai	Asp	Glu 150	Cys	
				140					147				•	130		
tac	atc	ccc	aac	tac	ctc	cta	ccc	aaq	act	cag	ccc	gag	ctg	cag	tgg	835
														Gln		
•			155					160					165			
														ctg		883
Ala	Trp			Met	Glu	Glu	Tyr 175		Ser	Ala	Cys	Leu 180		Leu	ATA	
		170					1/3					100		•		
gag	cac	aaa	caa	ctc	ata	gca	caq	cac	cta	acc	cac	gtc	ago	cag	acc	931
														Gln		
	185		•			190					195					

_		_		-	_		_					-	 cct Pro	-	979
							-		-	_	_	-	 tgg Trp 230	-	1027
	-							_	_			-	 gac Asp	-	1075
_	_						-	-			-		cag Gln		1123
_	-	-	_	-		-		-		-	-	_	aaa Lys		1171
		-		-	_			_	_		-	-	aaa Lys		1219
	-			-		_						-	gcc Ala 310		1267
-			-			-							aag Lys		1315
				_			_	_		-	Gly	_	aac Asn		1363
	-	-		-		-							aac Asn	-	1411
													gtg Val		1459
													gcc Ala 390		1507

-											
			agc Ser								1555
			ccc Pro								1603
			gac Asp								1651
			atg Met 445				_	_	-		1699
			gtg Val								1747
			gtg Val								1795
			aaa Lys								1843
			tgc Cys				_	_		_	1891
gcg Ala 520			gtg Val 525								1939
			atg Met								1987
			aat Asn								2035
			ttc Phe								2083

		Leu Cys Ala	att cat cat ctg Ile His His Leu 595	
	· · ·		gat ctg gtc aac Asp Leu Val Asn 610	
	tgt gtt tct gag Cys Val Ser Glu 620		acttaaa ccaggt at Me 62	t Ala Ala
			aac ctc tct gag Asn Leu Ser Glu 640	
	•	Pro Gly Ala	ccg aag ccc aaa Pro Lys Pro Lys 655	
-			gtg ctt cct ggc Val Leu Pro Gly 670	-
		-	ggg gag ccc gtc Gly Glu Pro Val	
	•		gcc tac gac cag Ala Tyr Asp Gln 705	-
	•		aac cac gcc gac Asn His Ala Asp 720	
		u Asp Thr Ser	ttt ggg ggc aac Phe Gly Gly Asn 735	
			ctc gaa cct ctc Leu Glu Pro Leu 750	
			a aag aaa cgt ccg / Lys Lys Arg Pro	

				-		-							aag Lys 785			2711
													ggc Gly	_		2759
				-						-			gca Ala			2807
													gca Ala		_	2855
				-		_	_		-			_	tca Ser			2903
											-		acc Thr 865		_	2951
	-			_	_								tac Tyr	_		2999
		-	_		_		-	_		-			tac Tyr			3047
											-		cac	-		3095
			-	-		_	-						tgg Trp			3143
		_	_										gtc Val 945	_	-	3191
-	-	_		-		_				-			ctt Leu		_	3239

-															
acg gtt	caa	gtc	ttc	tcg	gac	tcg	gag	tac	cag	ctt	ccg	tac	gtc	ctc	3287
Thr Val	Gln \	Val	Phe	Ser	Asp	Ser	Glu	Tvr	Gln	Leu	Pro	Tvr	Val	Leu	
965					970			- 7 -		975			, •		
505	'				570					913					
ggc tct	gcg	cac	cag	ggc	tgc	ctc	cct	ccg	ttc	ccg	gcg	gac	gtg	ttc	3335
Gly Ser	Alal	His (	Gln	Gly	Cys	Leu	Pro	Pro	Phe	Pro	Ala	Asp	Val	Phe	
980				985					990					995	
atg att	cca /	~ a a ·	t a c	aac	t 2.0	cta	200	ctc	220	224	~~~	200		<b>acc</b>	3383
-	_					_	-					-		-	3303
Met Ile	Pro		-	GIA	Tyr	Leu	Thr	Leu	Asn	Asn	GTÀ	Ser	GIn	Ala	
		1	000				1	.005				1	010		
gtg gga	cgt t	tca	tcc	ttt	tac	tqc	ctq	gaa	tat	ttc	cct	tct	caq	atq	3431
Val Gly						-	-	-					_	_	
, ur cr	_				- 3 -	_		Olu	1 y L	1110			Q111		
	Τ,	015				ı	.020				ذ	1025			
ctg aga	acg	ggc .	aac	aac	ttt	acc	ttc	agc	tac	acc	ttt	gag	gaa	gtg	3479
Leu Arg	Thr (	Gly i	Asn	Asn	P'ne	Thr	Phe	Ser	Tvr	Thr	Phe	Glu	Glu	Val	
-	1030	•				.035					040				
					•	.033				_	.040				
cct ttc	cac	agc .	agc	tac	gcg	cac	agc	cag	agc	ctg	gac	cgg	ctg	atg	3527
Pro Phe	His	Ser .	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu	Asp	Arg	Leu	Met	
1045	)			3	050				1	1055					
													·		
aat cct	ctc	atc	~ a c	<b>C</b> 2 2	* > 0	c+a	+ - +	<b>t</b> 2.0	c+~	226	202	3.C.t	~~~	224	3575
			_			-			-		-				3373
Asn Pro	Leu	iie.	Asp	GIn	Tyr	Leu	Tyr	Tyr	Leu	Asn	Arg	Thr			•
1069			1	.065				1	1070					1075	
cag tco	gga	agt	gcc	caa	aac	aag	gac	ttg	ctg	ttt	agc	cgt	ggg	tct	3623
Gln Ser	Glv	Ser	Ala	Gln	Asn	Lvs	Asp	Leu	Leu	Phe	Ser	Ara	Glv	Ser	
02 00.			080			_,_	_	1085				_	1090		
		_	000					.003				•	1090		
cca gct	ggc	atg	tct	gtt	cag	ccc	aaa	aac	tgg	cta	cct	gga	ccc	tgt	3671
Pro Ala	Gly	Met	Ser	Val	Gln	Pro	Lys	Asn	Trp	Leu	Pro	Gly	Pro	Cys	
	1	095				-	1100		_			1105			
	. –			•											
<b>.</b>					<b>L</b> = 4.										
tat cg		_	-	-						-				_	3719
Tyr Ar	g Gln	Gln	Arg	Val	Ser	Lys	Thr	Lys	Thr	Asp	Asn	Asn	Asn	Ser	
	1110					1115					1120				
aat tti	acc	taa	act	aat	act	tca	222	tat	aac	ctc	aat	gaa	cat	gaa	3767
					_								_		
Asn Pho		rrp	Inr	_		ser	гÀ2	Tyr			ASN	сту	Arg	GIU	
1125	•			-	1130					1135					
tcc ato	atc	aac	cct	gạc	act	gct	atg	gcc	tca	cac	aaa	gac	gac	gaa	3815
Ser Ile	lle i	Asn	Pro	Gly	Thr	Ala	Met	Ala	Ser	His	Lys	Asp	Asp	Glu	
1140				145					1150		•	•		1155	

gac aag ttc ttt Asp Lys Phe Phe					3863
gcc gga gct tca Ala Gly Ala Ser 1175			Val Met Ile		3911
gag gaa att aaa Glu Glu Ile Lys 1190	Ala Thr Asn		-		3959
gtg gca gtc aat Val Ala Val Asn 1205		-		22 2	4007
gtg cat gct atg Val His Ala Met 1220		Pro Gly Met			4055
gtg tac ctg cag Val Tyr Leu Gln					4103
cac ttt cac ccg His Phe His Pro 1255			Phe Gly Leu	-	4151
cct cct cag atc Pro Pro Gln Ile 1270	Leu Ile Lys	=		-	4199
gcg gag ttt tca Ala Glu Phe Ser 1285	-	<u>-</u>			4247
aca gga caa gtg Thr Gly Gln Val 1300		Ile Glu Trp		=	4295
agc aag cgc tgg Ser Lys Arg Trp				_	4343
tct gcc aac gtt Ser Ala Asn Val 1335	=		Asn Gly Leu		4391

cct cgc ccc att ggc acc cgt tac ctt acc cgt ccc ctg taattacgtg 4440

Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu

1350 1355 1360

ttaatcaata aaccggttga ttcgtttcag ttgaactttg gtctcctgtc cttcttatct 4500

tatcggttac catggttata gcttacacat taactgcttg gttgcgcttc gcgataaaag 4560
acttacgtca tcgggttacc cctagtgatg gagttgccca ctccctctct gcgcgctcgc 4620
tcgctcggtg gggcctgcgg accaaaggtc cgcagacggc agagctctgc tctgccggcc 4680
ccaccgagcg agcgagcgcg cagagaggga gtgggcaa 4718

<210> 2

<211> 623

<212> PRT

<213> AAV-1

<400> 2

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys . 160 Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Glr. Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245. Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys ·Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430

Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460

Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480

Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val
500 505 510

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 535 540

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 620

<210> 3

<211> 736

<212> PRT

<213> AAV-1

<400> 3

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly
145 150 155 160

Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175

Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190

Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205

Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220

Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu

-				245					250					255	
Tyr	Lys	Gln	Ile 260	Ser	Ser	Ala	Ser	Thr 265	Gly	Ala	Ser	Asn	Asp 270	Asņ	His
Туr	Phe	Gly 275	Tyr	Ser	Thr	Pro	Trp 280	Gly	Tyr	Phe	Asp	Phe 285	Asn	Arg	Phe
His	Cys 290	His	Phe	Ser	Pro	Arg 295	Asp	Trp	Gln	Arg	Leu 300	Ile	Asn	Asn	Asn
Trp 305	Gly	Phe	Arg	Pro	Lys 310	Arg	Leu	Asn	Phe	Lys 315	Leu	Phe	Asn	Ile	Gln 320
Val	Lys	Glu	Val	Thr 325	Thr	Asn	Asp	Gly	Val 330	Thr	Thr	Ile	Ala	Asn 335	Asn
Leu	Thr	Ser	Thr 340	Val	Gln	Val	Phe	Ser 345	Asp	Ser	Glu	Tyr	Gln 350	Leu	Pro
Tyŗ	Val	Leu 355	Gly	Ser	Ala	Hıs	Gln 360	Gly	Cys	Leu	Pro	Pro 365	Phe	Pro	Ala
Asp	Val 370	Phe	Met	Ile	Pro	Gln 375	Tyr	Gly	Tyr	Leu	Thr 380	Leu	Asn	Asn	Gly
ser 385	Gln	Ala	Val	Gly	Arg 390	Ser	Ser	Phe	Tyr	Cys 395	Leu	Glu	Tyr	Phe	Pro 400
Ser	Gln	Met	Leu	Arg 405	Thr	Gly	Asn	Asn	Phe 410	Thr	Phe	Ser	Tyr	Thr 415	Phe
Glu	Glu	Val	Pro 420	Phe	His	Ser	Ser	Tyr 425	Ala	His	Ser	Gln	Ser 430	Leu	Asp
Arg	Leu	Met 435	Asn		Leu	Ile	Asp 440	Gln	Туr	Leu	Туr	Tyr 445	Leu	Asn	Arg
Thr	Gln 450	Asn	Gln	Ser	Gly	Ser 455	Ala	Gln	Asn	Lys	Asp 460	Leu	Leu	Phe	Ser
Arg 465	Gly	Ser	Pro	Ala	Gly 470	Met	Ser	Val	Gln	Pro 475	Lys	Asn	Trp	Leu	Pro 480
Gly	Pro	Cys	Tyr	Arg 485	Gln	Gln	Arg	Val	Ser 490	Lys	Thr	Lys	Thr	Asp 495	Asn

Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn

510

500 505

Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys 515 520 525

Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 535 540

Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 555 560

Thr Asp Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg 565 570 575

Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590

Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln 595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 620

Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu 625 630 635 640

Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655

Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

<210> 4

<211> 1872

<212> DNA

<2-13> AAV-1

<220> <221> CDS

<222> (1)..(1869)

<400> 4

atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac 48

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp

1 5 10 15

gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag 96 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att 144
Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
35 40 45

gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg 192 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

gtc caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt 240.

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val
65 70 75 80

cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag 288 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att 336
Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile
100 105 110

agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg 384 Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432
Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly
130 135 140

aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag 480 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

act cag ccc gag ctg cag tgg gcg tgg act aac atg gag gag tat ata 528

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 age gee tgt ttg aac etg gee gag ege aaa egg ete gtg geg eag eac Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 ctg acc cac gtc agc cag acc cag gag cag aac aag gag aat ctg aac 624 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 ccc aat tct gac gcg cct gtc atc cgg tca aaa acc tcc gcg cgc tac Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220 atg gag ctg gtc ggg tgg ctg gtg gac cgg ggc atc acc tcc gag aag 720 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255 tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag 816 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 atc atg gcg ctg acc aaa tcc gcg ccc gac tac ctg gta ggc ccc gct 864 Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285 ccg ccc gcg gac att aaa acc aac cgc atc tac cgc atc ctg gag ctg 912 Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 aac ggc tac gaa cct gcc tac gcc ggc tcc gtc ttt ctc ggc tgg gcc 960 Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 320 cag aaa agg ttc ggg aag cgc aac acc atc tgg ctg ttt ggg ccg qcc Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 335 acc acg ggc aag acc aac atc gcg gaa gcc atc gcc cac gcc gtg ccc Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 tto tac ggc tgc gtc aac tgg acc aat gag aac ttt ccc ttc aat gat

P <u>h</u> e	Туr	Gly 355	Cys	Val	Asn	Trp	Thr 360	Asn	Glu	Asn	Phe	Pro 365	Phe	Asn	Asp	
												_	-	acġ Thr	_	1152
												-	_	gtg Val	-	1200
														ccc Pro 415		1248
														aac Asn	_	1296
														aaa Lys		1344
														aag Lys	_	1392
														gag Glu		1440
														ccc Pro 495	_	1488
														tca Ser		1536
											-		-	ttt Phe	_	1584
														cag Gln		1632
ctg	ttt	ccc	tgc	aag	aca	tgc	gag	aga	atg	aat	cag	aat	ttc	aac	att	1680

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560

tgc ttc acg cac ggg acg aga gac tgt tca gag tgc ttc ccc ggc gtg 1728 Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575

tca gaa tct caa ccg gtc gtc aga aag agg acg tat cgg aaa ctc tgt 1776 Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590

gcc att cat ctg ctg ggg cgg gct ccc gag att gct tgc tcg gcc 1824 Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605

tgc gat ctg gtc aac gtg gac ctg gat gac tgt gtt tct gag caa taa 1872 Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 620

<210> 5

<211> 623

<212> PRT

<213> AAV-1

<400> 5

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu
50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu

PCT/US99/25694

WO 00/28061

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala

_	3	7	0

375

380

Lys 385	Val	Val	Glu	Ser	Ala 390	Lys	Ala	Ile	Leu	Gly 395	Gly	Ser	Lys,	Val	Arg 400
Val	Asp	Gln	Lys	Cys 405	Lys	Ser	Ser _.	Ala	Gln 410	Ile	Asp	Pro	Thr	Pro 415	Val
Ile	Val	Thr	Ser 420	Asn	Thr	Asn	Met	Cys 425	Ala	Val	Ile	Asp	Gly 430	Asn	Ser
Thr	Thr	Phe 435	Glu	His	Gln	Gln	Pro 440	Leu	Gln	Asp	Arg	Met 445	Phe	Lys	Phe
Glu	Leu 450	Thr	Arg	Arg	Leu	Glu 455	His	Asp	Phe	Gly	Lys 460	Val	Thr	Lys	Gln
Glu 465	Val	Lys	Glu	Phe	Phe 470	Arg	Trp	Ala	Gln	Asp 475	His	Val	Thr	Glu	Val 480
Ala	His	Glu	Phe	Туг 485	Val	Arg	Lys	Gly	Gly 490	Ala	Asn	Lys	Arg	Pro 495	Ala
Pro	Asp	Asp	Ala 500	Asp	Lys	Ser	Glu	Pro 505	Lys	Arg	Ala	Cys	Pro 510	Ser	Val
Ala	Asp	Pro 515		Thr	Ser	Asp	Ala 520	Glu	Gly	Ala	Pro	Val 525	Asp	Phe	Ala
Asp	Arg 530	Tyr	Gln	Asn	Lys	Cys 535	Ser	Arg	His	Ala	Gly 540	Met	Leu	Gln	Met
Leu 545		Pro	Cys	Lys	Thr 550	-	Glu	Arg	Met	Asn 555		Asn	Phe	Asn	Ile 560
Cys	Phe	Thr	His	Gly 565		Arg	Asp	Cys	Ser 570		Cys	Phe	Pro	Gly 575	
Ser	Glu	ı Ser	Gln 580		Val	Val	Arg	Lys 585		Thr	Tyr	Arg	Lys 590		Cys
Ala	Ile	His 595		Leu	ı Leu	Gly	Arg 600		Pro	Glu	ı Ile	Ala 605		Ser	: Ala
Cys	Asp	Leu	ı Val	. Asr	val	Asp		Asp	Asp	Cys	Val		Glu	Glñ	ı

<210> 6 <211> 1641 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1638) <400> 6 atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac 48 Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 5 gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att 144 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 4 ū gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg 192 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60 gtc caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 70 cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 95 acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135 aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag 480 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys

145-	150	155	160
		aac atg gag gag ta Asn Met Glu Glu Ty	r Île
		a cgg ctc gtg gcg ca s Arg Leu Val Ala Gl 190	<del>-</del>
		g aac aag gag aat ct n Asn Lys Glu Asn Le 205	
		a aaa acc tcc gcg cg r Lys Thr Ser Ala Ar 220	·
		g ggc atc acc tcc ga g Gly Ile Thr Ser Gl 235	= =
		c atc tcc ttc aac gc r Ile Ser Phe Asn Al 0 25	a Ala
		t ctg gac aat gcc gg a Leu Asp Asn Ala Gl 270	<del>-</del>
		tac ctg gta ggc cc Tyr Leu Val Gly Pr 285	-
	=	c tac cgc atc ctg ga e Tyr Arg Ile Leu Gl 300	
		c gtc ttt ctc ggc tg r Val Phe Leu Gly Tr 315	•
		c tgg ctg ttt ggg cc e Trp Leu Phe Gly Pr 0 33	o Ala
		c atc gcc cac gcc gt a Ile Ala His Ala Va	<del>-</del>

**-** 340 345 350

ttc	tac	ggc	tgc	qtc	aac	taa	acc	aat	gag	aac	ttt	ccc	ttc	aat	gat	1104
			-	-	Asn											
	-	355	-			•	360					365			•	
•	,			•												
tac	atc	gac	aaσ	ato	gtg	atc	taa	t.aa	gag	gag	aac	aag	ato	aca	acc	1152
_	_	_	_	_	Val							-	_	-	-	1102
0,5	370		_,_		• • • •	375	110	110	Ozu	014	380	2,5			714	
	3.0					3.3					500					
aan	atc	ata	aaa	tcc	gcc	aaa	acc	att	ctc	aac	aac	age	aan	ata	cac	1200
					Ala	_	_			-		_	_		-	1200
385	•••	• • • •	Olu	Jer	390	БуЗ	ΛIα	116	Deu	395	Gry	261	Буз	Val	400	
505					390					3,7,3					400	
ata	aac	caa	aan	tac	aag	tca	tcc	acc	cad	at C	asc	CCC	200	CCC	ata	1248
	-		-	-	Lys	-		-	_		-					1240
VOI	vab	0111	пуэ	405	гуs	261	261	AIG	410	116	Asp	PIO	1111		Val	
				405					410					415		
	~+-							•								1006
	_				acc		-	_	-			•			-	1296
TTE	vaı	Thr		Asn	Thr	Asn	Met	_	АТа	val	ile	Asp	_	Asn	Ser	
			420					425					430			
						-										
					cag							_				1344
Thr	Thr		Glu	His	Gln	Gln	Pro	Leu	Gln	Asp	Arg	Met	Phe	Lys	Phe	
		435					440					445				
gaa	ctc	acc	cgc	cgt	ctg	gag	cat	gac	ttt	ggc	aag	gtg	aca	aag	cag	1392
Glu	Leu	Thr	Arg	Arg	Leu	Glu	His	Asp	Phe	Gly	Lys	Val	Thr	Lys	Gln	
	450					455					460					
													•			
gaa	gtc	aaa	gag	ttc	ttc	cgc	tyg	gcg	cag	gat	cac	gtg	acc	gag	gtg	1440
Glu	Val	Lys	Glu	Phe	Phe	Arg	Trp	Ala	Gln	Asp	His	Val	Thr	Glu	Val	
465					470		•			475					480	
gcq	cat	qaq	ttc	tac	gtc	aga	aaq	aat	gga	acc	aac	aaa	aga	ccc	acc	1488
					Val											
	•••-		• • • •	485		9	2,0	<b></b> ,	490			_,_		495		-
				`					150	•				.,,		
ccc	ast	a = c	aca	ast		3.00	~~~		224	666	~~~	t a c		+ 02	ato	1536
					aaa											1336
Pro	Asp	Asp		Asp	Lys	Ser	GIU		ьуs	Arg	Ala	Cys		ser	vaı	
			500					505					510			
					tca											1584
Ala	Asp	Pro	Ser	Thr	Ser	Asp	Ala	Glu	Gly	Ala	Pro	Val	Asp	Phe	Ala	
		515					520					525	•	•		
		•														
gac	agg	tat	ggc	tgc	cga	tgg	tta	tct	tcc	aga	ttg	gct	cga	gga	caa	1632
Asp	Arg	Tyr	Gly	Cys	Arg	Trp	Leu	Ser	Ser	Arg	Leu	Ala	Arg	Gly	Gln	

540

**_** 530 535

cct ctc tga 1641 Pro Leu

545

<210> 7

<211> 546

<212> PRT

<213> AAV-1

<400> 7

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp

1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 175

Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 190

Le $\overline{u}$  Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 205

- Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220
- Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 240
- Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255
- Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 270
- Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285
- Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300
- Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 315
- Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 . 335
- Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 350
- Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 365
- Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 370 375 380
- Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400
- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 455 450 Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 470 475 465 Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 505 500 Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 Asp Arg Tyr Gly Cys Arg Trp Leu Ser Ser Arg Leu Ala Arg Gly Gln 535 Pro Leu 545 <210> 8 <211> 1200 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1197) <400> 8 atg gag ctg gtc ggg tgg ctg gtg gac cgg ggc atc acc tcc gag aag Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 15 5 10 1 cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct 96 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 20 144 tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 35 40 atc atg gcg ctg acc aaa tcc gcg ccc gac tac ctg gta ggc ccc gct Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 60 . 55 50

_											
			aaa Lys 70					_		-	240
			gcc Ala							-	288
			aag Lys			-			_	-	336
			aac Asn								384
			aac Asn		-					-	432
			gtg Val 150						-	-	480
			gcc Ala				_	_		-	528
			aag Lys								576
			acc Thr							_	624
			cag Gln								672
			ctg Leu 230						_	-	720
			ttc Phe								768

		gag Glu														816
		gac Asp 275	-	-		_			_		-	-			_	864
		cca Pro		_		-		-		-	_		-		•	912
-	-	tac Tyr				-		-				_		_	-	960
_		ccc Pro	-	_		_		-	-		_					1008
_		acg Thr			_	_	-	-		- •	-					1056
	-	tct Ser 355		_	_	_	•	_		_					-	1104
		cat His						_				-	_	_	_	1152
		ctg Leu													taa	1200

<210> 9

<211> 399

<212> PRT

<213> AAV-1

<400> 9

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys

1 5 10 15

Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala

٠,	
~	•

25

30

Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 35 40 45

Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 50 55 60

Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 65 70 75 80

Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 85 90 95

Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 100 105 110

Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 115 120 125

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 130 135 140

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 145 150 155 160

Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 165 170 175

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 180 185 190

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 195 200 205

Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 210 215 220

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 225 230 235 240

Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val
245 250 255

Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 260 265 270

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val

PCT/US99/25694

275 280 285

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 290 295 300 .

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 305 310 315 320

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 325 330 335

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 340 345 350

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 355 360 365

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 370 375 380

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 385 390 395

<210> 10

<211> 969

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (1)..(966)

<220>

<221> misc_feature

<222> (943)..(944)

<223> minor splice site

<400> 10

atg gag ctg gtc ggg tgg ctg gtc ggg ggc atc acc tcc gag aag 48
Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys
1 5 10 15

cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct 96 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 20 25 30

tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag 144

Se <u>r</u>	Asn	Ser 35	Arg	Ser	Gln	Ile	Lys 40	Ala	Ala	Leu	Asp	Asn 45	Ala	Gly	Lys	
	-		_					ccc Pro								192
_		-						cgc Arg								240
								ggc Gly								288
-								acc Thr 105								336
								gaa Glu								384
								aat Asn								432
_								tgg Trp								480
								att Ile								528
	-		-	Cys				gcc Ala 185						Pro	gtg Val	576
	-						-	Cys	_				Gly		agc Ser	624
		Phe					Pro					Met			ttt Phe	672
gaa	ctc	acc	cgc	cgt	ctg	gag	cat	gac	ttt	ggc	aaç	gtg	aca	aag	cag	720

Gl <u>u</u> 225	Leu	Thr	Arg	Arg	Leu 230	Glu	His	Asp	Phe	Gly 235	Lys	Val	Thr	Lys	Gln 240	
				ttc Phe 245												768
				tac Tyr												816
				gat Asp											_	864
				acg Thr												912
				tgc Cys							-	-	-			960
	ctc Leu	tga							-						÷	969
<211 <212	0> 11 l> 32 2> PF B> AF	22 RT														
<400	)> 1]	l														
Met 1	Glu	Leu	Val	Gly 5	Trp	Leu	Val	Asp	Arg 10	Gly	Ile	Thr	Ser	Glu 15	Lys	
Gln	Trp	Ile	Gln 20	Glu	Asp	Gln	Ala	Ser 25	Tyr	Ile	Ser	Phe	Asn 30	Ala	Ala	
Ser	Asn	Ser 35	Arg	Ser	Gln	Ile	Lys 40	Ala	Ala	Leu	Asp	Asn 45	Ala	Gly	Lys	
Ile	Met 50	Ala	Leu	Thr	Lys	Ser 55	Ala	Pro	Asp	Tyr	Leu 60	Val	Gly	Pro	Ala	
Pro 65	Pro	Ala	Asp	Ile	Lys 70	Thr	Asn	Arg	Ile	Tyr 75	Arg	Ile	Leu	Glu	Leu 80	

Asm Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 85 90 95

- Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 100 105 110
- Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 115 120 125
- Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 130 135 140
- Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 145 150 155 160
- Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 165 170 175
- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 180 185 190
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 195 200 205
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 210 215 220
- Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 225 230 235 240
- Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 245 250 255
- Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 260 265 270
- Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 275 280 285
- Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 290 295 300
- Asp Arg Tyr Gly Cys Arg Trp Leu Ser Ser Arg Leu Ala Arg Gly Gln 305 310 315

Pro Leu

<210> 12 <211> 2211 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(2208) <400> 12 atg gct gcc gat ggt tat ctt cca gat tgg ctc gag gac aac ctc tct Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser 10 gag ggc att cgc gag tgg tgg gac ttg aaa cct gga gcc ccg aag ccc Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro aaa gcc aac cag caa aag cag gac gac ggc cgg ggt ctg gtg ctt cct 144 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 40 ggc tac aag tac ctc gga ccc ttc aac gga ctc gac aag ggg gag ccc 192 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 55 gtc aac gcg gcg gac gca gcg gcc ctc gag cac gac aag gcc tac gac Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 80 cag cag ctc aaa gcg ggt gac aat ccg tac ctg cgg tat aac cac gcc Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 gac gcc gag ttt cag gag cgt ctg caa gaa gat acg tct ttt ggg ggc Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly 100 105 aac ctc ggg cga gca gtc ttc cag gcc aag aag cgg gtt ctc qaa cct 384 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 125 ctc ggt ctg gtt gag gaa ggc gct aag acg gct cct gga aag aaa cqt 432 Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140 ccg gta gag cag tcg cca caa gag cca gac tcc tcc tcg ggc atc qqc

Pro- 145	Val	Glu	Gln	Ser	Pro 150	Gln	Glu	Pro	Asp	Ser 155	Ser	Ser	Gly	Ile	Gly 160	
				_		_	aaa Lys	_	_					_		528
							gat Asp									576
				_			cct Pro 200			_	-					624
-		-	-	-			gaa Glu		_	-					•	672
					-	_	tcc Ser			_		-	-	-		720
				-			gcc Ala	_								768
	_				_	-	tca Ser	_		_	_		-			816
				-			tgg Trp 280				_			_		864
-	-					-	gac Asp		-	-						912
					-	-	ctc Leu									960
-	_		-	_	-		gat Asp		-				-			1008
ctt	acc	agc	acg	gtt	caa	gtc	ttc	tcg	gac	tcg	gag	tac	cag	ctt	ccg	1056

Leu_	Thr	Ser	Thr 340	Val	Gln	Val	Phe	Ser 345	Asp	Ser	Glu	Tyr	Gln 350	Leu	Pro	
	Val		ggc Gly				_		-			_		_		1104
			atg Met													1152
			gtg Val								-	_				1200
			ctg Leu		_							-				1248
			cct Pro 420			_	-				-	_	_	-	-	1296
			aat Asn				_			-			-			1344
			cag Gln								-	_	-		_	1392
			cca Pro													1440
			tat Tyr											-		1488
			aat Asn 500													1536
			tcc Ser								_	-				1584
gac	gac	gaa	gac	aag	ttc	ttt	ccc	atg	agc	ggt	gtc	atg	att	ttt	gga	1632

Asp	Asp 530	Glu	Asp	Lys	Phe	Phe 535	Pro	Met	Ser	Gly	Val 540	Met	Ile	Phe	Gly	
	-	_	_		_				-	-	-	aat Asn	-	-		1680
												gcc Ala		_	-	1728
								_	-	_	_	aca	_			1776
												atg Met 605				1824
-	-	_			_	_					-	aaa Lys				1872
	-					_	,			-		ggc ggc				1920
-					-						_	cct Pro	•			1968
				-			-		-		-	tca Ser				2016
												tgg Trp 685		_	•	2064
									_		_	tac Tyr				2112
											-	aac Asn				2160
tat	act	gag	cct	cgc	ссс	att	ggc	acc	cgt	tac	ctt	acc	cgt	ccc	ctg	2208

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

<210> 13

<211> 736

<212> PRT

<213> AAV-1

<400> 13

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

1 10 15

Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly
145 150 155 160

Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190

Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205

- Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220
- Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240
- Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255
- Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270
- Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285
- His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300
- Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315
- Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335
- Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350
- Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365
- Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 380
- Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400
- Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe 405 410 415
- Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430
- Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg 435 440 445

Thr_Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His €15 Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu 725 730 <210> 14 <211> 1800 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1797) <400> 14 acg gct cct gga aag aaa cgt ccg gta gag cag tcg cca caa gag cca 48 Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro 10 gac too too tog ggc atc ggc aag aca ggc cag cag ccc gct aaa aag Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys 20 25 aga ctc aat ttt ggt cag act ggc gac tca gag tca gtc ccc gat cca 144 Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro 35 40 45 caa cet etc gga gaa eet eea gea ace eee get get gtg gga eet act 192 Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr aca atg gct tca ggc ggt ggc gca cca atg gca gac aat aac gaa qqc 240 Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly 65 70 75 gcc gac gga gtg ggt aat gcc tca gga aat tgg cat tgc gat tcc aca 288 Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr 85 90 tgg ctg ggc gac aga gtc atc acc acc agc acc cgc acc tgg gcc ttg Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu 100 110 ccc acc tac aat aac cac ctc tac aag caa atc tcc agt gct tca acg 384 Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr

120

125

115

					aac Asn										432
					aga Arg 150										480
_	_				aac Asn						-	-			528
					atc Ile	-	-	Glu	-	_	_		•		576
-				-	aat Asn			-	-	-		-		_	624
					ctt Leu						-		-		672
_			_		ccg Pro 230	 -			_		_				720
					aat Asn										768
	-	_	-		ttc Phe		-	-	-	-	•				816
					acc Thr		-					_	-		864
					ctg Leu										912
					aac Asn 310							-	-		960

		gac Asp				_					-		_		Val	1008
_		aaa Lys							-			-	-	•	-	1056
		aca Thr 355			-				-						, ,	1104
-		aaa Lys						-	-							1152
	-	atg Met	-				-	-	-	_	-				-	1200
		gtc Val													Thr	1248
-	-	gac Asp		-	_			-	-		-			-	act' Thr	1296
		gtg Val 435	_		_	_					-	_			-	1344
Ser	450	Ser		Asp	Pro	Ala 455	Thr	Gly	Asp	Val	460	Ala	Met	Gly	Ala	1392
		ggc											_		Pro 480	1440
		gcc Ala						-					•			1488
	_	ggc Gly					-		_			_				1536

-	•															
														gct		1584
Lys	Asn		Pro	Val	Pro	Ala		Pro	Pro	Ala	Glu		Ser	Ala	Thr	
•		515					520		•			525			•	
									_							
aag	ttt	gct	tca	ttc	atc	acc	caa	tac	tcc	aca	gga	caa	gtg	agt	gtg	1632
Lys	Phe	Ala	Ser	Phe	Ile	Thr	Gln	Tyr	Ser	Thr	Gly	Gln	Val	Ser	Val	
	530					535					540					
gaa	att	gaa	tgg	gag	ctq	cag	aaa	gaa	aac	agc	aag	cgc	tgg	aat	ccc	1680
Glu	Ile	Glu	Trp	Glu	Leu	Gln	Lys	Glu	Asn	Ser	Lys	Arg	Trp	Asn	Pro	
545			-		550		•			555	-	_	_		560	
gaa	ata	caq	tac	aca	tcc	aat	tat	αca	aaa	tct	αcc	aac	att	gat	ttt	1728
-		_						_					-	Asp		
014			- ] -	565			-,-	,,,,,	570					575		
				303					3.0					3,3		
201	ata	<b>~</b> ~ ~	336	225	993	~ F +	* > F	a.c.t	a 2 a	cct	cac		a + +	ggc	3.00	1776
		_									-			Gly		1770
THE	vaı	Asp		ASII	GIY	Leu	Tyr		Giu	PIO	Arg	FIO		GIY	1111	
			580					585					590			
																1000
_				-	ccc	-	taa									1800
Arg	Tyr		Thr	Arg	Pro	Leu									*	
		595				•										•
		_														
	0> 1															
<21	1> 5	99														

<212> PRT

<213> AAV-1

<400> 15

Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro 1 5 10 15

Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys 20 25 30

Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro 35 40 45

Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr 50 55 60

Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly 65 70 75 80

Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr

Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val 

Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val

**340 345 350** 

Ser Lys Thr Lys Thr Asp Asn Asn Ser Asn Phe Thr Trp Thr Gly 355 360 365

Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly 370 375 380

Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met 385 390 395 400

Ser Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr 405 410 415

Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Ile Lys Ala Thr 420 425 430

Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln 435 440 445

Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala 450 455 460

Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro 465 470 475 480

Ile Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro 485 490 495

Leu Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile 500 505 510

Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr 515 520 525

Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val 530 535 540

Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro 545 550 555 560

Glu Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe 565 570 575

Thr Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr 580 585 590

Arg Tyr Leu Thr Arg Pro Leu

- 595

<210> 16 <211> 1605 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1) ... (1602) <400> 16 atg gct tca ggc ggt ggc gca cca atg gca gac aat aac gaa ggc gcc Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala gac gga gtg ggt aat gcc tca gga aat tgg cat tgc gat tcc aca tgg Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp 20 25 ctg ggc gac aga gtc atc acc acc agc acc cgc acc tgg gcc ttg ccc Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro 35 45 acc tac aat aac cac ctc tac aag caa atc tcc agt gct tca acg ggg 192 Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly 50 55 gcc agc aac gac aac cac tac ttc ggc tac agc acc ccc tgg ggg tat 240 Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr 65 ttt gat ttc aac aga ttc cac tgc cac ttt tca cca cgt gac tgg cag 288 Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln 85 90 cga ctc atc aac aac tgg gga ttc cgg ccc aag aga ctc aac ttc 336 Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe 105 aaa ctc ttc aac atc caa gtc aag gag gtc acg acg aat gat ggc gtc 384 Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val 115 120 125 aca acc atc gct aat aac ctt acc agc acg gtt caa gtc ttc tcg gac 432 Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser Asp 130 135 140

•	-												
	gag Glu		Leu		-					-		_	480
	cct Pro											tac Tyr	528
	acg Thr												576
	ctg Leu												624
	ttc Phe 210												672
	agc Ser									-			720
	tat Tyr							-	-	-			768
	gac Asp										_	-	816
	aaa Lys										_		864
	aca Thr 290											-	912
	aaa Lys					-							960
	atg Met										-	-	1008

ggt gt Gly Va	-						_	_		-				-	1056
ttg ga Leu As		-	_			_	-		-			•			1104
cct gt Pro Va 37	l Ala		-	-				-	-	_			_	-	1152
agc ag Ser Se 385		_					_				_				1200
cct go					_	_	-			-	_				1248
tgg go															1296
atg gg Met G	gc ggc Ly Gly 435				-		_			-					1344
Asn T	eg cct nr Pro 50	-												_	1392
-	et tca la Ser											-		•	1440
_	aa tgg lu Trp		_	-		-		_	_	-				-	1488
	ag tac ln Tyr					-			_		_	-			1536
	ac aac sp Asn 515														1584

tac ctt acc cgt ccc ctg taa Tyr Leu Thr Arg Pro Leu 530 1605

<210> 17

<211> 534

<212> PRT

<213> AAV-1

<400> 17

Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala 1 5 10 15

Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp
20 25 30

Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro 35 40 45

Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly
50 55 60

Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr
65 70 75 80

Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln 85 90 95

Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe 100 105 110

Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val 115 120 125

Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser Asp 130 135 140

Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr
165 170 175

Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr 180 185 190

Cys_Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe 195 200 205

- Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr Ala 210 215 220
- His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr 225 230 235 240
- Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn 245 250 255
- Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val Gln 260 265 270
- Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser 275 280 285
- Lys Thr Lys Thr Asp Asn Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala 290 295 300
- Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr 305 310 315 320
- Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met Ser 325 330 335
- Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala 340 345 350
- Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn 355 360 365
- Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln Ser 370 380
- Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala Leu 385 390 395 400
- Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile 405 410 415
- Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro Leu
  420 425 430
- Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys 435 440 445

Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys
450 455 460

Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu 465 470 475 480

Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu
485 490 495

Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr 500 505 510

Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg
515 520 525

Tyr Leu Thr Arg Pro Leu 530

<210> 18

<211> 4681

<212> DNA

<213> aav-2

<400> 18

teggecacte cetetetege eyetegeteg etcactgagg cegggegace aaaggtegee 60 cgaegecegg getttgeeeg gyeggeetea gtgagegage gagegegaag agaggagtg 120 gecaacteca teactagggg teectggagg ggtggagteg tgaegtgaat taegteatag 180 ggttagggag gteetgtatt agaggteacg tgagtgttt gegacatttt gegacaceat 240 gtggteacge tgggtattta agecegagtg ageacgeagg gtetecattt tgaageggga 300 ggtttgaacg egeageegee atgeegggt tetacgagat tgtgattaag gteeceageg 360 acettgaegg geatetgeee ggeatttetg acagetttgt gaactgggtg geegagaagg 420 aatgggagt geegeagat tetgacatyg atetgaatet gattgageag geaceeetga 480 eegtggeega gaagetgeag egegacttee tgaeggaatg gegeegtgtg agtaaggeee 540 eggaggeeet tetettgtg caattegaga agggagagag etactteeae atgeaegge 600 tegtggaaac caeeggggtg aaateeatgg tettgggaeg tetteetgagt cagattegeg 720

tcacaaagac cagaaatggc gccggaggcg ggaacaaggt ggtggatgag tgctacatcc 780 ccaattactt gctccccaaa acccagccty agctccagtg ggcgtggact aatatggaac 840 agtatttaag cgcctgtttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga 900 cgcacgtgtc gcagacgcag gagcagaaca aagagaatca gaatcccaat tctgatgcgc 960 cggtgatcag atcaaaact tcagccaggt acatggagct ggtcgggtgg ctcgtggaca 1020 aggggattac ctcggagaag cagtggatcc aggaggacca ggcctcatac atctccttca 1080 atgcggcctc caactcgcgg tcccaaatca aggctgcctt ggacaatgcg ggaaagatta 1140 tgagcctgac taaaaccgcc cccqactacc tggtgggcca gcagcccgtg gaggacattt 1200 ccagcaatcg gatttataaa attttggaac taaacgggta cgatccccaa tatgcggctt 1260 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg 1320 ggcctgcaac taccgggaag accaacatcg cggaggccat agcccacact gtgcccttct 1380 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg 1440 tgatctggtg ggaggaggyg aagatgaccg ccaaggtcgt ggagtcggcc aaagccattc 1500 teggaggaag caaggtgege gtggaceaga aatgeaagte eteggeeeag atagaceega 1560 ctcccgtgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga 1620 ccttcgaaca ccagcagccg ttgcaagacc ggatgttcaa atttgaactc acccgccgtc 1680 tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cggtgggcaa 1740 aggatcacgt ggttgaggtg gagcatgaat tctacgtcaa aaagggtgga gccaagaaaa 1800 gacccgcccc cagtgacgca gatataagtg agcccaaacg ggtgcgcgag tcagttgcgc 1860 agccatcgac gtcagacgcg gaagcttcga tcaactacgc agacaggtac caaaacaaat 1920 gttctcgtca cgtgggcatg aatctgatgc tgtttccctg cagacaatgc gagagaatga 1980 atcagaattc aaatatctgc ttcactcacg gacagaaaga ctgtttagag tgctttcccg 2040 tgtcagaatc tcaaccggtt tctgtcgtca aaaaggcgta tcagaaactg tgctacattc 2100 atcatatcat gggaaaggtg ccagacgctt gcactgcctg cgatctggtc aatgtggatt 2160

tggatgactg catctttgaa caataaatga tttaaatcag gtatggctgc cgatggttat 2220 cttccagatt ggctcgagga cactctctct gaaggaataa gacagtggtg gaagctcaaa 2280 cctggcccac caccacaaa gcccgcagag cggcataagg acgacagcag gggtcttgtg 2340 cttcctgggt acaagtacct cggacccttc aacggactcg acaagggaga gccggtcaac 2400 gaggcagacg ccgcggccct cgagcacgac aaagcctacg accggcagct cgacagcgga 2460 gacaacccgt acctcaagta caaccacgcc gacgcggagt ttcaggagcg ccttaaagaa 2520 gatacgtett ttgggggeaa ceteggaega geagtettee aggegaaaaa gagggttett 2580 gaacctctcg gcctggttga ggaacctgtt aagacggctc cgggaaaaaa gaggccggta 2640 gagcactete etgtggagee agacteetee tegggaaceg gaaageeggg eeageageet 2700 gcaagaaaaa gattgaattt tggtcagact ggagacgcag actcagtacc tgacccccag 2760 cctctcggac agccaccagc agccccctct ggtctgggaa ctaatacgat ggctacaggc 2820 agtggcgcac caatggcaga caataacgag ggcgccgacg gagtgggtaa ttcctccgga 2880 aattggcatt gcgattccac atggatgggc gacagagtca tcaccaccag cacccgaacc 2940 tgggccctgc ccacctacaa caaccacctc tacaaacaaa tttccagcca atcaggagcc 3000 tegaaegaea ateaetaett tygetaeage acceettggg ggtattttga etteaaeaga 3060 ttccactgcc acttttcacc acgtgactgg caaagactca tcaacaacaa ctggggattc 3120 cgacccaaga gactcaactt caacctcttt aacattcaag tcaaagaggt cacgcagaat 3180 gacggtacga cgacgattgc caataacctt accagcacgg ttcaggtgtt tactgactcg 3240 gagtaccage tecegtacgt ceteggeteg gegeateaag gatgeeteee geegtteeea 3300 gcagacgtct tcatggtgcc acagtatgga tacctcaccc tgaacaacgg gagtcaggca 3360 gtaggacget etteatttta etgeetggag taettteett eteagatget gegtaeegga 3420 aacaacttta cetteageta caettttgag gaegtteett tecacageag etaegeteae 3480 agecagagte tggacegtet catgaateet eteategace agtacetgta ttaettgage 3540 agaacaaaca ctccaagtgg aaccaccacg cagtcaaggc ttcagttttc tcaggcccca 3600

gccagtgaca ttcgggacca gtctaggaac tggcttcctg gaccctgtta ccgccagcag 3660 cgagtatgaa agacatctgc ggataacaac aacagtgaat actcgtggac tggagctacc 3720 aagtaccace teaatggeag agaetetetg gtgaateegg ggeeegeeat ggeaageeae 3780 aaggacgatg aagaaaagtt ttttcctcag agcggggttc tcatctttgg gaagcaaggc 3840 tcagagaaaa caaatgtgaa cattgaaaag gtcatgatta cagacgaaga ggaaatccca 3900 acaaccaatc ccgtggctac ggagcagtat ggttctgtat ctaccaacct ccaqaqaqqc 3960 aacagacaag cagctaccgc agatgtcaac acacaaggcg ttcttccagg catggtctgg 4020 caggacagag atgtgtacct tcaggggccc atctgggcaa agattccaca cacggacgga 4080 cattttcacc cctctcccct catgggtgga ttcggactta aacaccctcc tccacagatt 4140 ctcatcaaga acaccccggt acctgcgaat ccttcgacca ccttcagtgc ggcaaagttt 4200 gcttccttca tcacacagta ctccacggga cacggtcagc gtggagatcg agtgggagct 4260 gcagaacgaa aacagcaaac gctggáatcc cgaaattcag tacacttcca actacaacaa 4320 gtctgttaat cgtggacttt accgtggata ctaatggcgt gtattcagag cctcgcccca 4380 ttggcaccag atacctgact cgtaatctgt aattgcttgt taatcaataa accgtttaat 4440 tegttteagt tgaactttgg tetetgegta tttettett atetagttte catggetaeg 4500 tagataagta gcatggcggg ttaatcatta actacaagga acccctagtg atggagttgg 4560 ccactccctc tctgcgcgct cgctcgctca ctgaggccgg gcgaccaaag gtcgcccgac 4620 gcccgggctt tgccccggcg gcctcagtga gcgagcgagc gcgcagagag ggagtgggca 4680 4681

<210> 19

<211> 4683

<212> DNA

<213> aav-6

<400> 19

ttggccactc cctctctgcg cgctcgctcg ctcactgagg ccgggcgacc aaaggtcgcc 60

cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc gagcgcgcag agagggagtg 120 gccaactcca tcactagggg ttcctggagg ggtggagtcg tgacgtgaat tacgtcatag 180 ggttagggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacaccat 240 gtggtcacgc tgggtattta agcccgagtg agcacgcagg gtctccattt tgaagcggga 300 ggtttgaacg cgcagcgcca tgccggggtt ttacgagatt gtgattaagg tccccagcga 360 ccttgacgag catctgcccg gcatttctga cagctttgtg aactgggtgg ccgagaagga 420 atgggagttg ccgccagatt ctgacatgga tctgaatctg attgagcagg cacccctgac 480 cgtggccgag aagctgcagc gcgacttcct ggtccactgg cgccgcgtga gtaaggcccc 540 ggaggccctc ttctttgttc agttcgagaa gggcgagtcc tacttccacc tccatattct 600 ggtggagacc acgggggtca aatccatggt gctgggccgc ttcctgagtc agattagcga 660 caagetggtg cagaceatet acegegggat egageegace etgeecaact ggttegeggt 720 gaccaagacg cgtaatggcg ccggaggggg gaacaaggtg gtggacgagt gctacatccc 780 caactacete etgeecaaga eteageeega getgeagtgg gegtggaeta acatggagga 840 gtatataage gegtgtttaa aeetggeega gegeaaaegg etegtggege aegaeetgae 900 ccacgtcagc cagacccagg agcagaacaa ggagaatctg aaccccaatt ctgacgcgcc 960 tgtcatccgg tcaaaaacct ccgcacgcta catggagctg gtcgggtggc tggtggaccg 1020 gggcatcace teegagaage agtggateea ggaggaeeag geetegtaca teteetteaa 1080 egeegeetee aastegeggt eeeagateaa ggeegetetg gacaatgeeg geaagateat 1140 ggcgctgacc aaatccgcgc ccgactacct ggtaggcccc gctccgcccg ccgacattaa 1200 aaccaaccgc atttaccgca tcctggagct gaacggctac gaccctgcct acgccggctc 1260 cgtctttctc ggctgggccc agaaaaggtt cggaaaacgc aacaccatct ggctgtttgg 1320 gccggccacc acgggcaaga ccaacatcgc ggaagccatc gcccacgccg tgcccttcta 1380 eggetgegte aactggacea atgagaactt teeetteaac gattgegteg acaagatggt 1440 gatctggtgg gaggagggca agatgacggc caaggtcgtg gagtccgcca aggccattct 1500

cggcggcagc aaggtgcgcg tggaccaaaa gtgcaagtcg tccgcccaga tcgatcccac 1560 ccccgtgatc gtcacctcca acaccaacat gtgcgccgtg attgacggga acagcaccac 1620 cttcgagcac cagcagccgt tgcaggaccy gatgttcaaa tttgaactca cccgccgtct 1680 ggagcatgac tttggcaagg tgacaaagca ggaagtcaaa gagttcttcc gctgggcgca 1740 ggatcacgtg accgaggtgg cgcatgagtt ctacgtcaga aagggtggag ccaacaacag 1800 accegeeec gatgaegegg ataaaagega geeeaagegg geetgeeeet eagtegegga 1860 tocatogacg toagacgogg aaggagotoc ggtggacttt gcogacaggt accaaaacaa 1920 atgttctcgt cacgcgggca tgcttcagat gctgtttccc tgcaaaacat gcgagagaat 1980 gaatcagaat ttcaacattt gcttcacgca cyggaccaga gactgttcag aatgtttccc 2040 cggcgtgtca gaatctcaac cggtcgtcag aaagaggacg tatcggaaac tctgtgccat 2100 teateatetg etggggeggg etecegagat tgettgeteg geetgegate tggteaacgt 2160 ggatctggat gactgtgttt ctgagcaata aatgacttaa accaggtatg gctgccgatg 2220 gttatettee agattggete gaggaeaace tetetgaggg catteggeag tggtgggaet 2280 tgaaacctgg agccccgaaa cccaaagcca accagcaaaa gcaggacgac ggccggggtc 2340 tggtgcttcc tggctacaag tacctcggac ccttcaacgg actcgacaag ggggagcccg 2400 tcaacgcggc ggatgcagcg gccctcgagc acgacaaggc ctacgaccag cagctcaaag 2460 egggtgacaa teegtacetg eggtataace acgeegaege egagttteag gagegtetge 2520 aagaagatac gtcttttggg ggcaacctcg ggcgagcagt cttccaggcc aagaagaggg 2580 ttctcgaacc ttttggtctg gttgaggaag gtgctaagac ggctcctgga aagaaacgtc 2640 cggtagagca gtcgccacaa gagccagact cctcctcggg cattggcaag acaggccagc 2700 agcccgctaa aaagagactc aattttggtc agactggcga ctcagagtca gtccccgacc 2760 cacaacctct cggagaacct ccagcaaccc ccgctgctgt gggacctact acaatggctt 2820 caggeggtgg egeaceaatg geagacaata aegaaggege egaeggagtg ggtaatgeet 2880 caggaaattg gcattgcgat tccacatggc tgggcgacag agtcatcacc accagcaccc 2940

gaacatgggc cttgcccacc tataacaacc acctctacaa gcaaatctcc agtgcttcaa 3000 cgggggccag caacgacaac cactacttcg gctacagcac cccctggggg tattttgatt 3060 tcaacagatt ccactgccat ttctcaccac gtgactggca gcgactcatc aacaacaatt 3120 ggggattecg geccaagaga eteaaettea agetetteaa eateeaagte aaggaggtea 3180 cgacgaatga tggcgtcacg accatcgcta ataaccttac cagcacggtt caagtcttgt 3240 cggactcgga gtaccagttc ccgtacgtcc tcggctctgc gcaccagggc tgcctcctc 3300 cgttcccggc ggacgtgttc atgattccgc agtacggcta cctaacgctc aacaatggca 3360 gccaggcagt gggacgctca tccttttact gcctggaata tttcccatcg cagatgctga 3420 gaacgggcaa taactttacc ttcagctaca ccttcgagga cgtgcctttc cacagcagct 3480 acgcgcacag ccagagcetg gaccggetga tgaateetet catcgaccag tacetgtatt 3540 acctgaacag aactcacaat cagtccygaa gtgcccaaaa caaggacttg ctgtttagcc 3600 gtgggtctcc agctggcatg tetgttcage ccaaaaactg getacetgga ccctgttace 3660 ggcagcagcg cgtttctaaa acaaaaacag acaacaacaa cagcaacttt acctggactg 3720 gtgcttcaaa atataacctt aatgggcgtg aatctataat caaccctggc actgctatgg 3780 cctcacacaa agacgacaaa gacaagttct ttcccatgag cggtgtcatg atttttggaa 3840 aggagagege eggagettea aacaetgeat tggacaatgt catgateaca gacgaagagg 3900 aaatcaaagc cactaacccc gtggccaccg aaagatttgg gactgtggca gtcaatctcc 3960 agagcagcag cacagaccct gcgaccggag atgtgcatgt tatgggagcc ttacctggaa 4020 tggtgtggca agacagagac gtatacctgc agggtcctat ttgggccaaa attcctcaca 4080 cggatggaca ctttcacccg tctcctctca tgggcggctt tggacttaag cacccgcctc 4140 ctcagatcct catcaaaaac acgcctgttc ctgcgaatcc tccggcagag ttttcggcta 4200 caaagtttgc ttcattcatc acccagtatt ccacaggaca agtgagcgtg gagattgaat 4260 gggagctgca gaaagaaaac aycaaacgct ggaatcccga agtgcagtat acatctaact 4320 atgcaaaatc tgccaacgtt gatttcactg tggacaacaa tggactttat actgagcctc 4380

<210> 20

<211> 16

<212> DNA

<213> rep binding motif

<400> 20

gctcgctcgc tcgctg

16

Jonal Application No

PCT/US 99/25694 A CLASSIFICATION OF SUBJECT MATTER IPC 7 -C12N15/86 C12N C12N15/35 C12N5/10 A61K48/00 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. RUTLEDGE E. A. ET AL.: "Infectious clones Y 1-23 and vectors derived from adeno-associated virus (AAV) serotypes other than AAV type 2." JOURNAL OF VIROLOGY, vol. 72, no. 1, January 1998 (1998-01), pages 309-319, XP002137089 ISSN: 0022-538X cited in the application the whole document Y WO 98 11244 A (SAFER BRIAN ;US HEALTH 1-23 (US); CHIORINI JOHN A (US); KOTIN ROBERT M) 19 March 1998 (1998-03-19) the whole document Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents: "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the International "X" document of particular relevance; the claimed invention filing date cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another "Y" document of particular relevance; the claimed invention citation or other special reason (as specified) cannot be considered to involve an inventive step when the document is combined with one or more other such docu-"O" document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled in the art. document published prior to the International filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 8 May 2000 22/05/2000 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijewijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,

Mandl, B

Fax: (+31-70) 340-3018

## INTERNATIONAL SE. RCH REPORT

Int Honal Application No PCT/US 99/25694

\(\(\text{O} = -1\)-	allan Bookini Cara	PCT/US 99/25694
etegory *	etion) DOCUMENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
, X	XIAO W. ET AL.: "Gene therapy vectors based on adeno-associated virus type 1." JOURNAL OF VIROLOGY, vol. 73, no. 5, May 1999 (1999-05), pages 3994-4003, XP002137090 ISSN: 0022-538X the whole document	1-23
	t	
	·	

...emational application No.

## INTERNATIONAL SEARCH REPORT

PCT/US 99/25694

Boxi	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This Inte	emational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:  Remark: Although claims 18-20 and 22, as far as an in vivo application is concerned, are directed to a method of treatment of the human or animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)
This Int	emational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remar	k on Protest  The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

information on patent family members

Int Honal Application No PCT/US 99/25694

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9811244 A	19-03-1998	AU 4645697 A EP 0932694 A	02-04-1998 04-08-1999